

11-08-00

A Box

SEA

DOCKET NO. : MOR-0003

PATENT

11/07/00



jc951 U.S. PTO

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In Re Application of:Nicholas C. Nicolaides, Luigi Grasso, and
Philip M. Sassjc951 U.S. PTO
09/707468**Serial No.:** Not assigned**Group Art Unit:** Not assigned**Filing Date:** November 7, 2000**Examiner:** Not assigned**For:** METHODS FOR GENERATING GENETICALLY ALTERED ANTIBODY-
PRODUCING CELL LINES WITH IMPROVED ANTIBODY
CHARACTERISTICS**EXPRESS MAIL LABEL NO:** EL568026574US
DATE OF DEPOSIT: November 7, 2000

Box ☒ Sequence
☐ Provisional ☐ Design

Assistant Commissioner for Patents
Washington DC 20231

Sir:

PATENT APPLICATION TRANSMITTAL LETTER

Transmitted herewith for filing, please find

☒ A Utility Patent Application under 37 C.F.R. 1.53(b).

It is a continuing application, as follows:

☐ continuation ☐ divisional ☐ continuation-in-part of prior application number
_____/_____.

☐ A Provisional Patent Application under 37 C.F.R. 1.53(c).
☐ A Design Patent Application (submitted in duplicate).

Including the following:

☐ Provisional Application Cover Sheet.

☒ New or Revised Specification, including pages 1 to 41 containing:

☒ Specification

☒ Claims

☒ Abstract

☐ Substitute Specification, including Claims and Abstract.

☐ The present application is a continuation application of Application No. _____ filed _____. The present application includes the Specification of the parent application which has been revised in accordance with the amendments filed in the parent application. Since none of those amendments incorporate new matter into the parent application, the present revised Specification also does not include new matter.

☐ The present application is a continuation application of Application No. _____ filed _____, which in turn is a continuation-in-part of Application No. _____ filed _____. The present application includes the Specification of the parent application which has been revised in accordance with the amendments filed in the parent application. Although the amendments in the parent C-I-P application may have incorporated new matter, since those are the only revisions included in the present application, the present application includes no new matter in relation to the parent application.

☐ A copy of earlier application Serial No. _____ Filed _____, including Specification, Claims and Abstract (pages 1 - @@), to which no new matter has been added TOGETHER WITH a copy of the executed oath or declaration for such earlier application and all drawings and appendices. Such earlier application is hereby incorporated into the present application by reference.

☐ Please enter the following amendment to the Specification under the Cross-Reference to Related Applications section (or create such a section) : "This Application:

☐ is a continuation of ☐ is a divisional of ☐ claims benefit of U.S. provisional Application Serial No. _____ filed _____

-
-
-
- ☐ Signed Statement attached deleting inventor(s) named in the prior application.
- ☐ A Preliminary Amendment.
- ☒ 7 Sheets of ☒ Formal ☐ Informal Drawings.
- ☐ Petition to Accept Photographic Drawings.
- ☐ Petition Fee
- ☒ An ☐ Executed ☒ Unexecuted Declaration or Oath and Power of Attorney.
- ☐ An Associate Power of Attorney.
- ☐ An ☐ Executed ☐ Copy of Executed Assignment of the Invention to _____
- ☐ A Recordation Form Cover Sheet.
- ☐ Recordation Fee - \$40.00.
- ☐ The prior application is assigned of record to _____
- ☐ Priority is claimed under 35 U.S.C. § 119 of Patent Application No. _____
filed _____ in _____ (country).
- ☐ A Certified Copy of each of the above applications for which priority is
claimed:
- ☐ is enclosed.
- ☐ has been filed in prior application Serial No. _____ filed _____.
- ☐ An ☐ Executed or ☐ Copy of Executed Earlier Statement Claiming Small Entity
Status under 37 C.F.R. 1.9 and 1.27
- ☐ is enclosed.
- ☐ has been filed in prior application Serial No. _____ filed _____,
said status is still proper and desired in present case.

- ☒ Diskette Containing DNA/Amino Acid Sequence Information.
- ☒ Statement to Support Submission of DNA/Amino Acid Sequence Information.
- ☐ The computer readable form in this application _____, is identical with that filed in Application Serial Number _____, filed _____. In accordance with 37 CFR 1.821(e), please use the ☐ first-filed, ☐ last-filed or ☐ only computer readable form filed in that application as the computer readable form for the instant application. It is understood that the Patent and Trademark Office will make the necessary change in application number and filing date for the computer readable form that will be used for the instant application. A paper copy of the Sequence Listing is ☐ included in the originally-filed specification of the instant application, ☐ included in a separately filed preliminary amendment for incorporation into the specification.
- ☐ Information Disclosure Statement.
- ☐ Attached Form 1449.
- ☐ Copies of each of the references listed on the attached Form PTO-1449 are enclosed herewith.
- ☐ A copy of Petition for Extension of Time as filed in the prior case.
- ☐ Appended Material as follows: _____.
- ☒ Return Receipt Postcard (should be specifically itemized).
- ☐ Other as follows: _____

_____.

FEE CALCULATION:

- ☐ Cancel in this application original claims _____ of the prior application before calculating the filing fee. (At least one original independent claim must be retained for filing purposes.)


			SMALL ENTITY		NOT SMALL ENTITY	
			RATE	FEE	RATE	FEE
PROVISIONAL APPLICATION			\$75.00	\$	\$150.00	\$
DESIGN APPLICATION			\$160.00	\$	\$320.00	\$
UTILITY APPLICATIONS BASE FEE			\$355.00	\$	\$710.00	\$710.00
UTILITY APPLICATION; ALL CLAIMS CALCULATED AFTER ENTRY OF ALL AMENDMENTS						
	No. Filed	No. Extra				
TOTAL CLAIMS	72 - 20 =	52	\$9 each	\$	\$18 each	\$936.00
INDEP. CLAIMS	7 - 3 =	4	\$40 each	\$	\$80 each	\$320.00
FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM			\$135	\$	\$270	\$
ADDITIONAL FILING FEE				\$		\$
TOTAL FILING FEE DUE				\$		\$1,966.00

- ☒ A Check is enclosed in the amount of \$ 1,966.00 .
- ☒ The Commissioner is authorized to charge payment of the following fees and to refund any overpayment associated with this communication or during the pendency of this application to deposit account 23-3050. This sheet is provided in duplicate.
- ☐ The foregoing amount due.
- ☒ Any additional filing fees required, including fees for the presentation of extra claims under 37 C.F.R. 1.16.
- ☒ Any additional patent application processing fees under 37 C.F.R. 1.17 or 1.20(d).
- ☐ The issue fee set in 37 C.F.R. 1.18 at the mailing of the Notice of Allowance.
- ☒ The Commissioner is hereby requested to grant an extension of time for the appropriate length of time, should one be necessary, in connection with this filing or any future filing submitted to the U.S. Patent and Trademark Office in the above-

identified application during the pendency of this application. The Commissioner is further authorized to charge any fees related to any such extension of time to deposit account 23-3050. This sheet is provided in duplicate.

SHOULD ANY DEFICIENCIES APPEAR with respect to this application, including deficiencies in payment of fees, missing parts of the application or otherwise, the United States Patent and Trademark Office is respectfully requested to promptly notify the undersigned.

Date: 11/7/00


Patrick J. Farley
Registration No. 42,524

Woodcock Washburn Kurtz
Mackiewicz & Norris LLP
One Liberty Place - 46th Floor
Philadelphia PA 19103
Telephone: (215) 568-3100
Facsimile: (215) 568-3439

METHODS FOR GENERATING GENETICALLY ALTERED ANTIBODY- PRODUCING CELL LINES WITH IMPROVED ANTIBODY CHARACTERISTICS

TECHNICAL FIELD OF THE INVENTION

5 The invention is related to the area of antibody maturation and cellular production. In particular, it is related to the field of mutagenesis.

BACKGROUND OF THE INVENTION

10 The use of antibodies to block the activity of foreign and/or endogenous polypeptides provides an effective and selective strategy for treating the underlying cause of disease. In particular is the use of monoclonal antibodies (MAb) as effective therapeutics such as the FDA approved ReoPro (Glaser, V. (1996) Can ReoPro repolish tarnished monoclonal therapeutics? *Nat. Biotechnol.* 14:1216-1217), an anti-platelet MAb from Centocor; Herceptin (Weiner, L.M. (1999) Monoclonal antibody therapy of cancer. *Semin. Oncol.* 26:43-51), an anti-Her2/neu MAb from Genentech; and Synagis (Saez-Llorens, X.E., *et al.* (1998) Safety and pharmacokinetics of an intramuscular humanized monoclonal antibody to respiratory syncytial virus in premature infants and infants with bronchopulmonary dysplasia. *Pediat. Infect. Dis. J.* 17:787-791), an anti-respiratory syncytial virus MAb produced by Medimmune.

20 Standard methods for generating MAbs against candidate protein targets are known by those skilled in the art. Briefly, rodents such as mice or rats are injected with a purified antigen in the presence of adjuvant to generate an immune response (Shield, C.F., *et al.* (1996) A cost-effective analysis of OKT3 induction therapy in cadaveric kidney

transplantation. *Am. J. Kidney Dis.* 27:855-864). Rodents with positive immune sera are sacrificed and splenocytes are isolated. Isolated splenocytes are fused to melanomas to produce immortalized cell lines that are then screened for antibody production. Positive lines are isolated and characterized for antibody production. The direct use of rodent MABs as

5 human therapeutic agents were confounded by the fact that human anti-rodent antibody (HARA) responses occurred in a significant number of patients treated with the rodent-derived antibody (Khazaeli, M.B., *et al.*, (1994) Human immune response to monoclonal antibodies. *J. Immunother.* 15:42-52). In order to circumvent the problem of HARA, the grafting of the complementarity determining regions (CDRs), which are the critical motifs
10 found within the heavy and light chain variable regions of the immunoglobulin (Ig) subunits making up the antigen binding domain, onto a human antibody backbone found these chimeric molecules are able to retain their binding activity to antigen while lacking the HARA response (Emery, S.C., and Harris, W.J. "Strategies for humanizing antibodies" In: ANTIBODY ENGINEERING C.A.K. Borrebaeck (Ed.) Oxford University Press, N.Y. 1995. pp.
15 159-183. A common problem that exists during the "humanization" of rodent-derived MABs (referred to hereon as HAb) is the loss of binding affinity due to conformational changes in the 3 dimensional structure of the CDR domain upon grafting onto the human Ig backbone (U.S. Patent No. 5,530,101 to Queen *et al.*). To overcome this problem, additional HAb vectors are usually needed to be engineered by inserting or deleting additional amino acid
20 residues within the framework region and/or within the CDR coding region itself in order to recreate high affinity HAbs (U.S. Patent No. 5,530,101 to Queen *et al.*). This process is a very time consuming procedure that involves the use of expensive computer modeling programs to predict changes that may lead to a high affinity HAb. In some instances the affinity of the HAb is never restored to that of the MAb, rendering them of little therapeutic
25 use.

Another problem that exists in antibody engineering is the generation of stable, high yielding producer cell lines that is required for manufacturing of the molecule for clinical materials. Several strategies have been adopted in standard practice by those skilled in the art to circumvent this problem. One method is the use of Chinese Hamster Ovary (CHO) cells
30 transfected with exogenous Ig fusion genes containing the grafted human light and heavy

chains to produce whole antibodies or single chain antibodies, which are a chimeric molecule containing both light and heavy chains that form an antigen-binding polypeptide (Reff, M.E. (1993) High-level production of recombinant immunoglobulins in mammalian cells. *Curr. Opin. Biotechnol.* 4:573-576). Another method employs the use of human lymphocytes derived from transgenic mice containing a human grafted immune system or transgenic mice containing a human Ig gene repertoire. Yet another method employs the use of monkeys to produce primate MAbs, which have been reported to lack a human anti-monkey response (Neuberger, M., and Gruggermann, M. (1997) Monoclonal antibodies. Mice perform a human repertoire. *Nature* 386:25-26). In all cases, the generation of a cell line that is capable of generating sufficient amounts of high affinity antibody poses a major limitation for producing sufficient materials for clinical studies. Because of these limitations, the utility of other recombinant systems such as plants are currently being explored as systems that will lead to the stable, high-level production of humanized antibodies (Fiedler, U., and Conrad, U. (1995) High-level production and long-term storage of engineered antibodies in transgenic tobacco seeds. *Bio/Technology* 13:1090-1093).

A method for generating diverse antibody sequences within the variable domain that results in HAbs and MAbs with high binding affinities to antigens would be useful for the creation of more potent therapeutic and diagnostic reagents respectively. Moreover, the generation of randomly altered nucleotide and polypeptide residues throughout an entire antibody molecule will result in new reagents that are less antigenic and/or have beneficial pharmacokinetic properties. The invention described herein is directed to the use of random genetic mutation throughout an antibody structure *in vivo* by blocking the endogenous mismatch repair (MMR) activity of a host cell producing immunoglobulins that encode biochemically active antibodies. The invention also relates to methods for repeated *in vivo* genetic alterations and selection for antibodies with enhanced binding and pharmacokinetic profiles.

In addition, the ability to develop genetically altered host cells that are capable of secreting increased amounts of antibody will also provide a valuable method for creating cell hosts for product development. The invention described herein is directed to the creation of genetically altered cell hosts with increased antibody production via the blockade of MMR.

The invention facilitates the generation of high affinity antibodies and the production of cell lines with elevated levels of antibody production. Other advantages of the present invention are described in the examples and figures described herein.

5 SUMMARY OF THE INVENTION

The invention provides methods for generating genetically altered antibodies (including single chain molecules) and antibody producing cell hosts *in vitro* and *in vivo*, whereby the antibody possess a desired biochemical property(s), such as, but not limited to, increased antigen binding, increased gene expression, and/or enhanced extracellular secretion
10 by the cell host. One method for identifying antibodies with increased binding activity or cells with increased antibody production is through the screening of MMR defective antibody producing cell clones that produce molecules with enhanced binding properties or clones that have been genetically altered to produce enhanced amounts of antibody product.

The antibody producing cells suitable for use in the invention include, but are not
15 limited to rodent, primate, or human hybridomas or lymphoblastoids; mammalian cells transfected and expressing exogenous Ig subunits or chimeric single chain molecules; plant cells, yeast or bacteria transfected and expressing exogenous Ig subunits or chimeric single chain molecules.

Thus, the invention provides methods for making hypermutable antibody-producing cells by introducing a polynucleotide comprising a dominant negative allele of a mismatch repair gene into cells that are capable of producing antibodies. The cells that are capable of producing antibodies include cells that naturally produce antibodies, and cells that are engineered to produce antibodies through the introduction of immunoglobulin encoding sequences. Conveniently, the introduction of polynucleotide sequences into cells is accomplished by transfection.

The invention also provides methods of making hypermutable antibody producing cells by introducing a dominant negative mismatch repair (MMR) gene such as *PMS2* (preferably human *PMS2*), *MLH1*, *PMS1*, *MSH2*, or *MSH2* into cells that are capable of producing antibodies. The dominant negative allele of a mismatch repair gene may be a truncation mutation of a mismatch repair gene (preferably a truncation mutation at codon 134,

or a thymidine at nucleotide 424 of wild-type *PMS2*). The invention also provides methods in which mismatch repair gene activity is suppressed. This may be accomplished, for example, using antisense molecules directed against the mismatch repair gene or transcripts.

Other embodiments of the invention provide methods for making a hypermutable antibody producing cells by introducing a polynucleotide comprising a dominant negative allele of a mismatch repair gene into fertilized eggs of animals. These methods may also include subsequently implanting the eggs into pseudo-pregnant females whereby the fertilized eggs develop into a mature transgenic animal. The mismatch repair genes may include, for example, *PMS2* (preferably human *PMS2*), *MLH1*, *PMS1*, *MSH2*, or *MSH2*. The dominant negative allele of a mismatch repair gene may be a truncation mutation of a mismatch repair gene (preferably a truncation mutation at codon 134, or a thymidine at nucleotide 424 of wild-type *PMS2*).

The invention further provides homogeneous compositions of cultured, hypermutable, mammalian cells that are capable of producing antibodies and contain a dominant negative allele of a mismatch repair gene. The mismatch repair genes may include, for example, *PMS2* (preferably human *PMS2*), *MLH1*, *PMS1*, *MSH2*, or *MSH2*. The dominant negative allele of a mismatch repair gene may be a truncation mutation of a mismatch repair gene (preferably a truncation mutation at codon 134, or a thymidine at nucleotide 424 of wild-type *PMS2*). The cells of the culture may contain *PMS2*, (preferably human *PMS2*), *MLH1*, or *PMS1*; or express a human *mutL* homolog, or the first 133 amino acids of hPMS2.

The invention further provides methods for generating a mutation in an immunoglobulin gene of interest by culturing an immunoglobulin producing cell selected for an immunoglobulin of interest wherein the cell contains a dominant negative allele of a mismatch repair gene. The properties of the immunoglobulin produced from the cells can be assayed to ascertain whether the immunoglobulin gene harbors a mutation. The assay may be directed to analyzing a polynucleotide encoding the immunoglobulin, or may be directed to the immunoglobulin polypeptide itself.

The invention also provides methods for generating a mutation in a gene affecting antibody production in an antibody-producing cell by culturing the cell expressing a dominant negative allele of a mismatch repair gene, and testing the cell to determine whether the cell

harbors mutations within the gene of interest, such that a new biochemical feature (*e.g.*, over-expression and/or secretion of immunoglobulin products) is generated. The testing may include analysis of the steady state expression of the immunoglobulin gene of interest, and/or analysis of the amount of secreted protein encoded by the immunoglobulin gene of interest. The invention also embraces prokaryotic and eukaryotic transgenic cells made by this process, including cells from rodents, non-human primates and humans.

Other aspects of the invention encompass methods of reversibly altering the hypermutability of an antibody producing cell, in which an inducible vector containing a dominant negative allele of a mismatch repair gene operably linked to an inducible promoter is introduced into an antibody-producing cell. The cell is treated with an inducing agent to express the dominant negative mismatch repair gene (which can be *PMS2* (preferably human *PMS2*), *MLH1*, or *PMS1*). Alternatively, the cell may be induced to express a human *mutL* homolog or the first 133 amino acids of hPMS2. In another embodiment, the cells may be rendered capable of producing antibodies by co-transfecting a preselected immunoglobulin gene of interest. The immunoglobulin genes of the hypermutable cells, or the proteins produced by these methods may be analyzed for desired properties, and induction may be stopped such that the genetic stability of the host cell is restored.

The invention also embraces methods of producing genetically altered antibodies by transfecting a polynucleotide encoding an immunoglobulin protein into a cell containing a dominant negative mismatch repair gene (either naturally or in which the dominant negative mismatch repair gene was introduced into the cell), culturing the cell to allow the immunoglobulin gene to become mutated and produce a mutant immunoglobulin, screening for a desirable property of said mutant immunoglobulin protein, isolating the polynucleotide molecule encoding the selected mutant immunoglobulin possessing the desired property, and transfecting said mutant polynucleotide into a genetically stable cell, such that the mutant antibody is consistently produced without further genetic alteration. The dominant negative mismatch repair gene may be *PMS2* (preferably human *PMS2*), *MLH1*, or *PMS1*. Alternatively, the cell may express a human *mutL* homolog or the first 133 amino acids of hPMS2.

The invention further provides methods for generating genetically altered cell lines that express enhanced amounts of an antigen binding polypeptide. These antigen-binding polypeptides may be, for example, immunoglobulins. The methods of the invention also include methods for generating genetically altered cell lines that secrete enhanced amounts of an antigen binding polypeptide. The cell lines are rendered hypermutable by dominant negative mismatch repair genes that provide an enhanced rate of genetic hypermutation in a cell producing antigen-binding polypeptides such as antibodies. Such cells include, but are not limited to hybridomas. Expression of enhanced amounts of antigen binding polypeptides may be through enhanced transcription or translation of the polynucleotides encoding the antigen binding polypeptides, or through the enhanced secretion of the antigen binding polypeptides, for example.

Methods are also provided for creating genetically altered antibodies *in vivo* by blocking the MMR activity of the cell host, or by transfecting genes encoding for immunoglobulin in a MMR defective cell host.

Antibodies with increased binding properties to an antigen due to genetic changes within the variable domain are provided in methods of the invention that block endogenous MMR of the cell host. Antibodies with increased binding properties to an antigen due to genetic changes within the CDR regions within the light and/or heavy chains are also provided in methods of the invention that block endogenous MMR of the cell host.

The invention provides methods of creating genetically altered antibodies in MMR defective Ab producer cell lines with enhanced pharmacokinetic properties in host organisms including but not limited to rodents, primates, and man.

These and other aspects of the invention are provided by one or more of the embodiments described below. In one embodiment of the invention, a method for making an antibody producing cell line hypermutable is provided. A polynucleotide encoding a dominant negative allele of a MMR gene is introduced into an antibody-producing cell. The cell becomes hypermutable as a result of the introduction of the gene.

In another embodiment of the invention, a method is provided for introducing a mutation into an endogenous gene encoding for an immunoglobulin polypeptide or a single chain antibody. A polynucleotide encoding a dominant negative allele of a MMR gene is

introduced into a cell. The cell becomes hypermutable as a result of the introduction and expression of the MMR gene allele. The cell further comprises an immunoglobulin gene of interest. The cell is grown and tested to determine whether the gene encoding for an immunoglobulin or a single chain antibody of interest harbors a mutation. In another aspect of the invention, the gene encoding the mutated immunoglobulin polypeptide or single chain antibody may be isolated and expressed in a genetically stable cell. In a preferred embodiment, the mutated antibody is screened for at least one desirable property such as, but not limited to, enhanced binding characteristics.

In another embodiment of the invention, a gene or set of genes encoding for Ig light and heavy chains or a combination therein are introduced into a mammalian cell host that is MMR defective. The cell is grown, and clones are analyzed for antibodies with enhanced binding characteristics.

In another embodiment of the invention, a method will be provided for producing new phenotypes of a cell. A polynucleotide encoding a dominant negative allele of a MMR gene is introduced into a cell. The cell becomes hypermutable as a result of the introduction of the gene. The cell is grown. The cell is tested for the expression of new phenotypes where the phenotype is enhanced secretion of a polypeptide.

These and other embodiments of the invention provide the art with methods that can generate enhanced mutability in cells and animals as well as providing cells and animals harboring potentially useful mutations for the large-scale production of high affinity antibodies with beneficial pharmacokinetic profiles.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. Hybridoma cells stably expressing PMS2 and PMS134 MMR genes.

Shown is steady state mRNA expression of MMR genes transfected into a murine hybridoma cell line. Stable expression was found after 3 months of continuous growth. The (-) lanes represent negative controls where no reverse transcriptase was added, and the (+) lanes represent samples reverse transcribed and PCR amplified for the MMR genes and an internal housekeeping gene as a control.

Figure 2. Creation of genetically hypermutable hybridoma cells. Dominant negative

MMR gene alleles were expressed in cells expressing a MMR-sensitive reporter gene. Dominant negative alleles such as PMS134 and the expression of MMR genes from other species results in antibody producer cells with a hypermutable phenotype that can be used to produce genetically altered immunoglobulin genes with enhanced biochemical features as well as lines with increased Ig expression and/or secretion. Values shown represent the amount of converted CPRG substrate which is reflective of the amount of function β -galactosidase contained within the cell from genetic alterations within the pCAR-OF reporter gene. Higher amounts of β -galactosidase activity reflect a higher mutation rate due to defective MMR.

Figure 3. Screening method for identifying antibody-producing cells containing antibodies with increased binding activity and/or increased expression/secretion

Figure 4. Generation of a genetically altered antibody with an increased binding activity. Shown are ELISA values from 96-well plates, screened for antibodies specific to hIgE. Two clones with a high binding value were found in HB134 cultures.

Figure 5. Sequence alteration within variable chain of an antibody (a mutation within the light chain variable region in MMR-defective HB134 antibody producer cells). Arrows indicate the nucleotide at which a mutation occurred in a subset of cells from a clone derived from HB134 cells. Panel A: The change results in a Thr to Ser change within the light chain variable region. The coding sequence is in the antisense direction. Panel B: The change results in a Pro to His change within the light chain variable region.

Figure 6. Generation of MMR-defective clones with enhanced steady state Ig protein levels. A Western blot of heavy chain immunoglobulins from HB134 clones with high levels of MAb (>500ngs/ml) within the conditioned medium shows that a subset of clones express higher steady state levels of immunoglobulins (Ig). The H36 cell line was used as a control to measure steady state levels in the parental strain. Lane 1: fibroblast cells (negative control); Lane 2: H36 cell; Lane 3: HB134 clone with elevated MAb levels; Lane 4: HB134 clone with elevated MAb levels; Lane 5: HB134 clone with elevated MAb levels.

Methods have been discovered for developing hypermutable antibody-producing cells by taking advantage of the conserved mismatch repair (MMR) process of host cells.

Dominant negative alleles of such genes, when introduced into cells or transgenic animals,

increase the rate of spontaneous mutations by reducing the effectiveness of DNA repair and thereby render the cells or animals hypermutable. Hypermutable cells or animals can then be utilized to develop new mutations in a gene of interest. Blocking MMR in antibody-producing cells such as but not limited to: hybridomas; mammalian cells transfected with genes encoding for Ig light and heavy chains; mammalian cells transfected with genes encoding for single chain antibodies; eukaryotic cells transfected with Ig genes, can enhance the rate of mutation within these cells leading to clones that have enhanced antibody production and/or cells containing genetically altered antibodies with enhanced biochemical properties such as increased antigen binding. The process of MMR, also called mismatch proofreading, is carried out by protein complexes in cells ranging from bacteria to mammalian cells. A MMR gene is a gene that encodes for one of the proteins of such a mismatch repair complex. Although not wanting to be bound by any particular theory of mechanism of action, a MMR complex is believed to detect distortions of the DNA helix resulting from non-complementary pairing of nucleotide bases. The non-complementary base on the newer DNA strand is excised, and the excised base is replaced with the appropriate base, which is complementary to the older DNA strand. In this way, cells eliminate many mutations that occur as a result of mistakes in DNA replication.

Dominant negative alleles cause a MMR defective phenotype even in the presence of a wild-type allele in the same cell. An example of a dominant negative allele of a MMR gene is the human gene *hPMS2-134*, which carries a truncating mutation at codon 134 (SEQ ID NO:15). The mutation causes the product of this gene to abnormally terminate at the position of the 134th amino acid, resulting in a shortened polypeptide containing the N-terminal 133 amino acids. Such a mutation causes an increase in the rate of mutations, which accumulate in cells after DNA replication. Expression of a dominant negative allele of a mismatch repair gene results in impairment of mismatch repair activity, even in the presence of the wild-type allele. Any allele which produces such effect can be used in this invention. Dominant negative alleles of a MMR gene can be obtained from the cells of humans, animals, yeast, bacteria, or other organisms. Such alleles can be identified by screening cells for defective MMR activity. Cells from animals or humans with cancer can be screened for defective mismatch repair. Cells from colon cancer patients may be particularly useful. Genomic

DNA, cDNA, or mRNA from any cell encoding a MMR protein can be analyzed for variations from the wild type sequence. Dominant negative alleles of a MMR gene can also be created artificially, for example, by producing variants of the *hPMS2-134* allele or other MMR genes. Various techniques of site-directed mutagenesis can be used. The suitability of such alleles, whether natural or artificial, for use in generating hypermutable cells or animals can be evaluated by testing the mismatch repair activity caused by the allele in the presence of one or more wild-type alleles, to determine if it is a dominant negative allele.

A cell or an animal into which a dominant negative allele of a mismatch repair gene has been introduced will become hypermutable. This means that the spontaneous mutation rate of such cells or animals is elevated compared to cells or animals without such alleles. The degree of elevation of the spontaneous mutation rate can be at least 2-fold, 5-fold, 10-fold, 20-fold, 50-fold, 100-fold, 200-fold, 500-fold, or 1000-fold that of the normal cell or animal. The use of chemical mutagens such as but limited to methane sulfonate, dimethyl sulfonate, O6-methyl benzadine, MNU, ENU, etc. can be used in MMR defective cells to increase the rates an additional 10 to 100 fold that of the MMR deficiency itself.

According to one aspect of the invention, a polynucleotide encoding for a dominant negative form of a MMR protein is introduced into a cell. The gene can be any dominant negative allele encoding a protein, which is part of a MMR complex, for example, *PMS2*, *PMS1*, *MLH1*, or *MSH2*. The dominant negative allele can be naturally occurring or made in the laboratory. The polynucleotide can be in the form of genomic DNA, cDNA, RNA, or a chemically synthesized polynucleotide.

The polynucleotide can be cloned into an expression vector containing a constitutively active promoter segment (such as but not limited to CMV, SV40, Elongation Factor or LTR sequences) or to inducible promoter sequences such as the steroid inducible pIND vector (Invitrogen), where the expression of the dominant negative MMR gene can be regulated. The polynucleotide can be introduced into the cell by transfection.

According to another aspect of the invention, an immunoglobulin (Ig) gene, a set of Ig genes or a chimeric gene containing whole or parts of an Ig gene can be transfected into MMR deficient cell hosts, the cell is grown and screened for clones containing genetically altered Ig genes with new biochemical features. MMR defective cells may be of human,

primates, mammals, rodent, plant, yeast or of the prokaryotic kingdom. The mutated gene encoding the Ig with new biochemical features may be isolated from the respective clones and introduced into genetically stable cells (*i.e.*, cells with normal MMR) to provide clones that consistently produce Ig with the new biochemical features. The method of isolating the

5 Ig gene encoding Ig with new biochemical features may be any method known in the art. Introduction of the isolated polynucleotide encoding the Ig with new biochemical features may also be performed using any method known in the art, including, but not limited to transfection of an expression vector containing the polynucleotide encoding the Ig with new biochemical features. As an alternative to transfecting an Ig gene, a set of Ig genes or a

10 chimeric gene containing whole or parts of an Ig gene into an MMR deficient host cell, such Ig genes may be transfected simultaneously with a gene encoding a dominant negative mismatch repair gene into a genetically stable cell to render the cell hypermutable.

Transfection is any process whereby a polynucleotide is introduced into a cell. The process of transfection can be carried out in a living animal, *e.g.*, using a vector for gene

15 therapy, or it can be carried out *in vitro*, *e.g.*, using a suspension of one or more isolated cells in culture. The cell can be any type of eukaryotic cell, including, for example, cells isolated from humans or other primates, mammals or other vertebrates, invertebrates, and single celled organisms such as protozoa, yeast, or bacteria.

In general, transfection will be carried out using a suspension of cells, or a single cell,

20 but other methods can also be applied as long as a sufficient fraction of the treated cells or tissue incorporates the polynucleotide so as to allow transfected cells to be grown and utilized. The protein product of the polynucleotide may be transiently or stably expressed in the cell. Techniques for transfection are well known. Available techniques for introducing polynucleotides include but are not limited to electroporation, transduction, cell fusion, the

25 use of calcium chloride, and packaging of the polynucleotide together with lipid for fusion with the cells of interest. Once a cell has been transfected with the MMR gene, the cell can be grown and reproduced in culture. If the transfection is stable, such that the gene is expressed at a consistent level for many cell generations, then a cell line results.

An isolated cell is a cell obtained from a tissue of humans or animals by mechanically

30 separating out individual cells and transferring them to a suitable cell culture medium, either

with or without pretreatment of the tissue with enzymes, *e.g.*, collagenase or trypsin. Such isolated cells are typically cultured in the absence of other types of cells. Cells selected for the introduction of a dominant negative allele of a mismatch repair gene may be derived from a eukaryotic organism in the form of a primary cell culture or an immortalized cell line, or may be derived from suspensions of single-celled organisms.

A polynucleotide encoding for a dominant negative form of a MMR protein can be introduced into the genome of an animal by producing a transgenic animal. The animal can be any species for which suitable techniques are available to produce transgenic animals. For example, transgenic animals can be prepared from domestic livestock, *e.g.*, bovine, swine, sheep, goats, horses, etc.; from animals used for the production of recombinant proteins, *e.g.*, bovine, swine, or goats that express a recombinant polypeptide in their milk; or experimental animals for research or product testing, *e.g.*, mice, rats, guinea pigs, hamsters, rabbits, etc. Cell lines that are determined to be MMR defective can then be used as a source for producing genetically altered immunoglobulin genes *in vitro* by introducing whole, intact immunoglobulin genes and/or chimeric genes encoding for single chain antibodies into MMR defective cells from any tissue of the MMR defective animal.

Once a transfected cell line or a colony of transgenic animals has been produced, it can be used to generate new mutations in one or more gene(s) of interest. A gene of interest can be any gene naturally possessed by the cell line or transgenic animal or introduced into the cell line or transgenic animal. An advantage of using such cells or animals to induce mutations is that the cell or animal need not be exposed to mutagenic chemicals or radiation, which may have secondary harmful effects, both on the object of the exposure and on the workers. However, chemical mutagens may be used in combination with MMR deficiency, which renders such mutagens less toxic due to an undetermined mechanism. Hypermutable animals can then be bred and selected for those producing genetically variable B-cells that may be isolated and cloned to identify new cell lines that are useful for producing genetically variable cells. Once a new trait is identified, the dominant negative MMR gene allele can be removed by directly knocking out the allele by technologies used by those skilled in the art or by breeding to mates lacking the dominant negative allele to select for offspring with a desired trait and a stable genome. Another alternative is to use a CRE-LOX expression

system, whereby the dominant negative allele is spliced from the animal genome once an animal containing a genetically diverse immunoglobulin profile has been established. Yet another alternative is the use of inducible vectors such as the steroid induced pIND (Invitrogen) or pMAM (Clonetech) vectors which express exogenous genes in the presence of corticosteroids.

Mutations can be detected by analyzing for alterations in the genotype of the cells or animals, for example by examining the sequence of genomic DNA, cDNA, messenger RNA, or amino acids associated with the gene of interest. Mutations can also be detected by screening for the production of antibody titers. A mutant polypeptide can be detected by identifying alterations in electrophoretic mobility, spectroscopic properties, or other physical or structural characteristics of a protein encoded by a mutant gene. One can also screen for altered function of the protein *in situ*, in isolated form, or in model systems. One can screen for alteration of any property of the cell or animal associated with the function of the gene of interest, such as but not limited to Ig secretion.

Examples of mismatch repair proteins and nucleic acid sequences include the following:

PMS2 (mouse) (SEQ ID NO:5)

```

MEQTEGVSTE CAKAIKPIDG KSVHQICSGQ VILSLSTAVK ELIENSVDAG ATTIDRLRKD 60
YGVDLIEVSD NGCGVEEENF EGLALKHHST KIQEFADLTQ VETFGFRGEA LSSLCALSDV 120
TISTCHGSAS VGTRLVFDHN GKITQKTPYP RPKGTTVSQV HLFYTLFVRV KEFQRNIKE 180
YSKMVQVLQA YCIISAGVRV SCTNQLGQGG RHAVVCTSGT SGMKENIGSV FGGKQLQSLI 240
FFVQLPPSDA VCEEYGLSTS GRHKTFSTFR ASFHSARTAP GGVOQTGSGS SSIRGPVTTQ 300
RSLSLSMRFY HMYNRHQYPF VVLNVSDVSE CVDINVTDPK RQILLQEEKL LLAVLKTSLI 360
25 GMFSDSANKL NVNQPLLDV EGNLVKLHTA ELEKPVPGKQ DNSPSLKSTA DEKRVASISR 420
LREAFSLHPT KEIKSRGPET AELTRSPFSE KRGVLSYSPS DVISYRGLRG SQDKLVSPDT 480
SPGDCMDREK IEKDSGLSST SAGSEEFST PEVASSFSDD YNVSSLEDRP SQETINCDDL 540
DCRPPTGTGS LKPEDHGYQC KALPLARLSP TNAKRFKTEE RPSNVNISQR LPPGPSTSA 600
EVDVAIKNNK RIVLLEFSLT SLAKRMKQLQ HLKQNKHEL SYRKFRACIK FGENQAAED 660
30 LRKEKISMF AEMEILQGQFN LGFIVTKLKE DLFLVDQHAA DEKYNFEMQL QHTVLAQORL 720
ITPQTNLNTA VNEAVLIENL EIFRKNGFDF VIDEADPCTAC TACTATGTAT KNTWFGPQDI 780
DELI FMLS DS FGVMCRPSRV RQMFASRACR KSMVIGTALN ASEMKKLITH MGEMDHPWNC 840
PHGRPTMRHV ANLDVISON 859

```

PMS2 (mouse cDNA) (SEQ ID NO:6)

```

gaattccggt gaaggtctctg aagaatttcc agattctctga gtatcattgg aggagacaga 60
taacctgtcg taggttaacg atggtgtata tgcaacacgaa atgggtgttc ctggagacgc 120
gtctttttccc gagagcggca ccgcaactct cccgcggtga ctgtgactgg aggagctctg 180
catccatgga gcaaacccgaa ggcgtgagta cagaatgtgc taaggccattg aagcctattg 240
atgggaagtc atgcatcaaa atttgttctg ggcaggtgat actcagttta agcaccgctg 300
tgaaggagtt gatagaaaaa atgtgtagatg ctggtgctac tactattgat ctaaggctta 360
aagactatgg ggtggacctc attgaagttt cagacaatgg atgtggggta gaagaagaaa 420

```

	actttgaagg	tctagctctg	aaacatcaca	catctaagat	tcaagagttt	gccgacctca	480
	cgagggttga	aacttttcggc	tttcgggggg	aagctctgag	ctctctgtgt	gcaactaagt	540
	atgtcacctat	atctacctgc	cacgggtctg	caagcgttgg	gactcgtactg	gtgtttgacc	600
	ataatgggaa	aatcacccag	aaaactccct	acccccgacc	taaaaggacc	acagtcaagt	660
5	gtgcagcaatt	atttttatca	ctaccogtgc	gttacaaaaga	gttttcagagg	aacattaaaa	720
	aggagatttct	caaaaagggtg	caggtctttac	aggcgtactg	tatcatctca	gcaggcgctc	780
	gtgtgaagct	cactaatcac	ctcggacagg	ggaagcggca	cgcgtgtggtg	tgcaacaagg	840
	gcacgtctcgg	catgaaggaa	aatatcgggt	ctgtgttttg	ccagaagcag	gtgtcaaaagg	900
	tcattctcttt	tgtttcagctg	ccccctagtg	acgctgtgtg	tgaaagattac	ggcctgagca	960
10	ctctcaggagc	ccacaaaaac	ttttctacgt	ttcggccttc	atttcacagt	gcacgcacgg	1020
	cgccggggagg	agtgcaaacg	acaggcagtt	ttttctcact	aatcagaagg	cctgtgacc	1080
	agcaaaaggtc	tctaagcttg	tcaatgaggt	tttatcacat	gtataaccgg	catcagttac	1140
	cattttgcgt	ccttaacggt	tcogttgact	cagaatgtgt	ggatattaat	gtacctccag	1200
	ataaaaggca	aattctacta	caagaagaga	agctatttgt	ggcgttttta	gaacacctct	1260
15	tgatagagct	gtttgacagt	gatgcacaaa	agcttaatgt	caaccagcag	caactgtcag	1320
	atgttgaagg	taacttagta	aagctgcata	ctgcagaact	agaaaagcct	gtgccagaaa	1380
	acgaagataa	ctctccttca	ctgaagagca	cagcagacga	gaaaagggtg	gaactccatt	1440
	ccagcgtcag	agaggccttt	ctctctcact	ctactaaaga	galcaagtct	aggggtccag	1500
20	agactgctga	actgcacagg	agttttccaa	gtgagaaaag	gggctgtgta	tcctcttacc	1560
	cttcagacag	catctctttac	agaggcctcc	gtggctcaga	ggacaaattg	gtgactccca	1620
	gcgaacagccc	tggtgactgt	atggacagag	agaaaaatga	aaaaagactca	agggctcaga	1680
	cgcactcagc	tggtcttgag	gaagagttca	gcaccccaga	agtgggcagt	gccttagcca	1740
	gtgactataa	cgtgagctcc	ctagaagaca	gacctcttga	ggaaaccata	aactgttggtg	1800
	actgggactc	cgcgtctcca	ggtagcaggac	agctcttgaa	gcacagaagac	catggatatac	1860
25	aatgcaaaag	cttaccttca	gctcgtctgt	cacccacaaa	tgccaaagcgc	ttcaagaagc	1920
	agggaaagacc	ctcaaatgtc	aaacatttgc	ttgttctctc	gtgtctcagc	agcacactgc	1980
	cagctgaggtg	cgatgtagcc	ataaaaaatg	ataagagaat	cgtgtcctct	gagttctctc	2040
	tgatttctct	agtcagacga	atgaagcagt	tacagcacct	aaaggcgcag	acaaaactac	2100
30	aactcagttta	cagaaaaatt	agggccaaga	tttgccctgg	agaaaaccac	gcagcagaag	2160
	atgaactctg	aaaaagagatt	agtaaatoga	gttttgacga	gatggagatc	ttgggtcagt	2220
	tttaactgttg	atttatagta	acccaaactga	aagaggacct	cttctcgtgt	gaccagcagt	2280
	ctcgggtaga	gaagtacaa	tttgagatgc	cacggtagca	cacggtgctc	gagccgcaga	2340
	ggctcatcac	acccagact	ctgaacttaa	ctgctgtcaa	tgagagctga	ctgatagaaa	2400
	atctggaaa	atccagaaag	aatggctttg	actttgtcat	tgataggatt	gctccagttca	2460
35	ctgaaagggc	taaatgtatt	tccttaccaa	ctagtaaaaa	ctggaccttt	ggaccccag	2520
	atatagatga	actgactctt	atgttaagtg	acagccctgg	gtctcatgtc	agccctccac	2580
	gagtcagaca	gatgtttgtc	tcagagacct	gtcggaaagt	agtcatgatt	ggaacgcggc	2640
	tcgaatgcgag	cagatgaag	aagctcatca	cccacatggg	tgagatggac	acccctgtgc	2700
40	actgccccca	cggcagtgcca	accatgaggg	acgttgcaca	ctctggaatc	atctctcaga	2760
	actgacacac	ccottgttagc	atagaagttta	ttacagattg	ttcggctttg	aaagagaagg	2820
	tttttaagtaa	cttgattacc	gttgatacaa	aattagcatg	ctgctctta	gtactgtgat	2880
	cttttaaaag	cagtggttaa	gcaggcatga	tggaagtttc	ctctagctca	gctactgtgg	2940
	tgatccggtg	ggagctcatg	tgagcccagg	actttgagac	cactccgagc	cacattcatg	3000
45	agactcaatt	caaggacaaa	aaaaaaaaa	tatttttgaa	gccttttaaa	aaaaaa	3056

PMS2 (human) (SEQ ID NO:7)

	MERAESSSTE	PAKAIKPIDR	KSVHQICSQG	VVLSLSTAVK	ELVENSIDAG	ATNIDLKLKD	60
	YGVDLLEIVSD	NGCGVEEENF	EGLTLKHHTS	KIQEFADLTQ	VETFGPRGEA	LSSLCLALSDV	120
	TIPTCHASAK	VGTRLMFDHN	GKIIQKTPYP	RPRGTTVSVQ	TLSTFLPVKH	KFQRNIKKE	180
50	YARMYQVLHA	YCIISAGIRV	SCTNQLGGQG	RQPVVCTGGS	PSIKENIGSV	FGGKQLQSLI	240
	PFYQLPPSDS	VCEEYGLSCS	DALHNLFYIS	GFISQCTHGV	GRSSDRQFF	FINNRPDCPA	300
	KVCRVLVNEVY	HMYNRHQYPF	VVLNLSVDSE	CVDINVTDPK	RQLLLQEEKL	LILAVLITSLI	360
	GMFDSVNNKL	NVSQQLLDV	EGNLLKMHA	DLEKPMVEKC	DQSPSLRTGE	EKKDVISIRL	420
55	REFSLRHTT	ENKPHSKPTP	EPRRSPLGQK	RGMLSSSTSG	AISDKGLVRP	QKEAVSSSHG	480
	PSPTDRAEV	EKDSGHGSTS	VDSGFSIPD	TGSHCSSEYA	ASSPGDRGSG	EHVDSQEKAP	540
	ETDSDSPVD	CHSNQEDTGC	KFRVLPQPTN	LATPNTKRFK	KEELISSSDI	CQKLNVTDQM	600
	SASQVDAVVK	INKKVQPLDF	SMSSLAKRIK	QLHHEAQOSE	GEQNYRKFFA	KICPGENAGA	660
	EDELRLKISK	TMPFAMEIIG	QFNLGFIITK	LNEDIFIVDQ	HATDEKYNEF	MLQHTVLQGG	720
	QRILIAQPTLN	LTVANEVLII	ENLEIFRKNL	FDPFVIDENAP	VTERAKLVIS	PTSKNVNPGP	780
60	QDVELIIFML	SDSPGVMCRP	SRVKQMFASR	ACRKSVMIGT	ALNTSEMKKL	ITHMGMDHP	840
	WNPCFGRPTM	RHIANLGVIS	QN				862

PMS2 (human cDNA) (SEQ ID NO:8)

cgaagccgat cgggtgttgc atccatggag cgagctgaga gctcagatgac agaacctgct 60
 aaggccatca aacctattga tcggaagtca gctccatcaga ttgtgctctgg gcaagtggtga 120
 5 ctgaggtctcaa gcaactgcggt aaaggaggtta gtgaaataca gctcggatgac tgggtgccact 180
 aatattgatac taagctctaa ggactattgga gtgagatctta ttgaaggtttc agacaatgga 240
 tgtggggtagt aagaagaaaaa cttcgaaggc ttaactctga aacatcacac atctaaagtat 300
 caaagattttg ccgacctaac tcagggttga aacttttggc ttcggggggga agctctgagc 360
 tcaatttggc cactgagcga tgtcaccatt tctacctgcc acgcatacggc gaaggttggca 420
 10 actcgaactga tttgtgatca caatgggaaa attatccaga aaaaaccctca cccctgccc 480
 agaggggacca cagtcagcgt gcagcagttta ttttccacac tacctgtgag ccaataaggaa 540
 tttcaaaagga atattaagaa ggagtatgcc aaaaatggtcc aggttctaca tgcatactgt 600
 atcattttcag caggcatccg tgtaatgtgc accaatcagc ttggacaagg aaaaacgacag 660
 cctgtggtat gcacaggtgg aagcccacgc ataaaggaaa atatcggtcc tgtgtttggg 720
 15 cagaagcagt tgcagaagcct cacttctttt gttcagctgc cccctagtga ctcagttgtg 780
 gaagagctgc gttttgagctg ttccgatgct ctgcataatc ttttttacaat ctcaggtttg 840
 atttcacaat gcacgcagct agtttgaagg agttcaacag acagacagtt tttctttatc 900
 aaccggcgcc cttgtgaccc agcaaaaggtc tgcagactcg tgaatgaggt caccacacatg 960
 tataatcgac accagtatcc atttgttgtt cttaacattt ctgttgattc agaattgcgtt 1020
 20 gatattcaat ttaactccaga taaaaggaaa attttgtctac aattgcttgc aggtttgttg 1080
 gcaatttttaa agactctctt gataggaatg tttgatagtg atgtcaacaa gctaaatgtc 1140
 agtcacagca cactctggtga ttttgaaggt aacttaataa aatgcgatgc agcggtattg 1200
 gaaaagccca ttgttgaaaa cgaggatcaa tccccttcat taaggactgg agaagaaaaa 1260
 25 aaagcagctg ccaattccag actcgcagag gcctttttct tctgcacac acagagaaac 1320
 aagcctcaga gcccaaaagc tccagaaaca agaaggagcc tctgacaaa ggcgtcctgag 1380
 atgctgtctt ctagcacttc aggtgccatc tctgacaaag gacctacgg ggtggagaag 1440
 gaggcagtag gttccagtcac cggaccagat cctcggtgat tctgaggggt tccagcatccc 1500
 actcgggggc acggcagcgc ttccgtggat tccggcaggg acagggggctc gcaggaaact 1560
 30 agtcaactga cgacgagtag tgcggccagc gactgaaact tttcagatgt ggaactgcct 1620
 gtggactctc agggagaaagc gactgaaact tttcagatgt ggaactgcct 1680
 tcaaacccgc aagatcacgg atgtaaaatt cgagtttttg ctagccaaac taactctgca 1740
 accccaacaa caaagcgctt taaaaaaaga gaaattcttt ccaagtctga ctaattgtcaa 1800
 aagttagtaa atactcagga catgtcagcc tctcaggttg atgtagctg gaaaattaat 1860
 35 aagaaagtgt tgcacctgga cttttctatg agttctttag cttaacgaat aagcagbta 1920
 catcatgaag cacagcaaa gaaagggaaa cagaattaca ggaagtttag ggcaaaagatt 1980
 tgtcctggag tggaaatcat agccgaagat gaactaaaga aagagataag taaaacgatg 2040
 ttgcagaaa tggaaatcat tggtcagttt aacctgggag ttataataac caaactgaat 2100
 40 gaggatatct ccatagtgcc ccagcatgcc acggcagaga agtataactt cgaagtgtctg 2160
 cagcagcaca ccgtgtccca ggggcagagc ctcatagcac ctacgactct caacttaact 2220
 ctgtgttaag aagctgttct cccagatcat ctgaaaatat ttagaagaag tggctttgat 2280
 ttgtttatcg atgaaaatgc tccagtcatc gaaagggtca aactgattc ctgtcccaact 2340
 agtataaaat ggaactctcg accccaggag gtccgatgac tgcattctac gctgagcgac 2400
 agccctgggg tcatgtgccg gccctcccca gtcaagcaga gtatgaaaga catgagcctgc 2460
 45 ccgaagtcgg tgatgattgc gactgctctt aacacagaag agatgaagaa actgatcacc 2520
 ccatgggggg agatggacca cccctggaac tgcgccatgc gaaggccaac catgagacac 2580
 ttgcgcaacc tgggtgtcat ttctcagaac tgaccgtagt caactgtatg aataatttgt 2640
 attatcgcag atttttatgt ttgaaagac agagtcttca cttaaccttt ttgtttttaa 2700
 50 atgaaacctg ctacttaaaa aaaaatacaca tcaaccccat ttaaagtgat tcttgagaa 2760
 cttttcaaac c 2771

PMS1 (human) (SEQ ID NO:9)

MKQLFAATVR LSSSQITTS VVSVVKELIE NSLDAGATSV DVKLENYGFD KIEVRDNGEG 60
 IKVADPAPMA KMYYSKINS HEDLENLTYY GERGEALGSI CTAIEVLITP RTAADNFSTQ 120
 55 YVLDGSGHIL SQPKSHLGGQ TTVTALRLFK NLPVRKQFYS TAKKCKDEIK KIQDLMSFG 180
 ILKPDRLRIV VHNKAVTWQK SRVSDHKML MSVLIRAVNM NMESFOYHSE ESYQILSGFL 240
 FKCDADHSFT SLSTPERSFI FINSRPVHKL NLKCLKESTR LVBPFFLKID 300
 VPTADVDVNL TPDKQVLLQ NKESVLIALE NLMTTCYGLP PSTNSYENNK TDSVDAADIVL 360
 SKTAEDTVLF NKVESSGKNY SNVDTSVIPF QNDHMDCLNGH RTDGDGFGYGH 420
 CSSEISNDIK NTKNAFQDIS MSNVSWNSVG TSEYKTCFIS VKHQTQSENG NKQHDISEGE 480
 60 NEEAGLENS SEESADEWSR GNILKNSVGE NLEPQKILVF EKSLPCKVSN NTPPIPEQNM 540
 LNEDSCNKKK NVIDNKSGKV TAYDLLSNRV IKKPMASAL FVQDHRFPQL IENPKTSLED 600

ATLQIEELWK	TLSEEEKLKY	E EKATKD LER	YNSQMKRAIE	QESQMSLKD G	RKKIKPTSAW	660
NLAQKHKKKT	SLSNPKLDE	LLQSQIEKR R	SNQIKMVQIP	FSMKNLKINF	KKQNKV DLEE	720
KDPECLINH L	RFPDAMLTS	KTEVM LNPY	RVEALLFRK	LLENHKLPAE	PLEKIMITE	780
SLFNGSHYLD	VLYKMTADDQ	RYSGSTYLS D	PRLTANGFKI	KLIPGV SITE	NYLEIEGMAN	840
CLPFYGVADL	KEILNAILNR	NAKEVYECRP	RKVISYLEGE	AVRLSRQLFM	YLSKEDIQDI	900
IYRMKHQFGN	EKECVHGRP	FFHLLTYLPE	TT			932

PMS1 (human) (SEQ ID NO:10)

10	ggcacgagtg	gctgcttgcg	gctagtggat	ggtaattgcc	tgccctcgcc	tagcagcaag	60
	ctgctctgtt	aaagcgcaaa	atgaacaat	tgccctcgcc	aacagttcga	ctccttccaa	120
	gttctcagat	catcacttcg	gtgggtcagtg	tgttaaaaga	gcttattgaa	aaactcctgg	180
	atgctgggtc	cacacgcgta	gatgttaaac	tgggagaacta	tggatttgat	aaaattggagg	240
	tgcgagataa	cggggaggggt	atcaaggctg	tgtatgcacc	tgtaatggca	atgaagtaact	300
	acacctcaaa	aataaatagt	catgaagatc	ttagaaattt	gacaaacttac	gggtttctgtg	360
15	gagaagcctt	ggggctcaatt	tgttgatag	ctgaggtttt	aattacaaca	agaacggcgtg	420
	ctgataattt	tagcacccag	tatgttttag	atggcagtg	ccactactct	ttccagaaac	480
	cttcacatct	tgttcaagggt	acaactgtaa	ctgctttaag	attatttaag	aatctacactg	540
	taagaaagca	gtttttactca	actgcacaaa	atgttaaaaa	tgaataaaaa	aagatccagag	600
	atctcctcat	gagcttttgtt	atccttaaac	ctgacttaag	gattgtcttt	gtacataaca	660
20	aggcagttat	ttggcagaaa	agcagagtat	cagatcacaa	gatggctctc	atgtcagttc	720
	tggggactgc	tgttatgaac	aatatggaat	cctttcagta	ccactctgaa	gaatctcaga	780
	tttatctcag	tggattttctt	ccaaagtgtg	atgcagacca	ctctttcact	agcttttccaa	840
	caccagaaa	aagtttcac	ttcataaaca	gtcgaccagt	actcaaaaa	gtatctctaa	900
	agtttaaccc	aatctatcac	aatctgaaat	gcotaaagga	atcatctcgt	ttgtattcgt	960
25	tttctcttct	gaaatctgat	gttctcaacg	ctgatgttga	tgtaaattta	acaccagata	1020
	aaagcccaat	attattacaa	aatgaagaa	ctgttttaaa	tgcttttgaa	atgtctatga	1080
	cgacttgata	tggacattta	cttagtcaaa	attcttatga	aaataataaa	acagatgttt	1140
	ccgcagctgc	catcgtctctt	agtaaaaaag	cagaaaacaga	tgtgcttttt	ataaagttgg	1200
	aatcatctgg	aaagaattat	tcaaatgttg	atacttcagt	catlccattc	caaaatgata	1260
30	tgcataatgt	tgaactctgga	aaaaaacctg	atgattgttt	aaatcaccaag	ataagtattg	1320
	gtgactttgg	ttatggtcat	tgtagttagt	aaatttctaa	cattgtataa	aaacctaaga	1380
	atgcatttca	ggacatttca	atgagtaagt	tatcatggga	gaactctcag	coggaatata	1440
	gtaaaacttg	ttttataatg	tcogttaagc	acacccagtc	agaaaattggc	acgaagaacc	1500
	atatagatga	tgaatggggaa	aatgaggaag	aagcagggtc	tggaaaactc	tcogaaatttt	1560
35	ctgcagatga	gtggagcagc	ggaaaataac	ttaaaaattc	agtgggagag	aatatttgac	1620
	ctgtgaaaa	tttagtgcct	gaaaaaagtt	taccatgtaa	agtaagtaat	ataattatct	1680
	caatccctga	acaaaatgat	cttaatgaag	attcatgtaa	caaaaaatca	aatgttaatg	1740
	ataataaatc	tggaagaatt	acagcttagt	atttacttag	caatcgagta	atacgaagac	1800
	coactgcacg	aagtgtctct	tttgttcaag	atcatcgctc	tcagtttctc	atagaaaatc	1860
40	ctaagactag	tttagaggat	gcaacacatc	aaattgaaga	actgtggaag	acattgagtg	1920
	aagaggaaaa	actgaaatat	gaagagaagg	ctactaaaga	cttggaacga	tacataagtc	1980
	aaatgaagag	agccatttga	caggagtcac	aaatgtcact	aaaaagatgg	agaaaaagaa	2040
	taaaaaccac	cagcgcatgg	aaatttggcc	agaagcacaa	gttaaaaaac	tcattatcta	2100
	atcaaccaaa	actgatgaga	ctccttcagt	cccaaaattg	aaaaaagagg	agtcacaaata	2160
45	tttaaaatg	acagatcccc	ttttctatga	aaaaacttaa	aaaaaatttt	aatgacacaaa	2220
	acaaagttga	cttagagacat	aaggatgaac	cttgcttgat	ccacaatact	aggtttctgt	2280
	atgatgtgct	aatgacatcc	aaaaacagag	taattgttat	aaatccatat	agagttaagc	2340
	agggcctgct	attttaaaaga	cttcttgaga	atcataaact	tctctcagag	ccactggaaa	2400
	agccaaattat	gtttaacagag	agtcttttta	atgatctcga	ttttttagac	gtttttataa	2460
50	aaatgacagc	agatgaccaa	agatcacagt	gatcaactta	ctctgtctgat	ctctgtcttta	2520
	cagcgcaattg	tttcaagata	aaatttgata	caggagtttt	aatctactgaa	ataactcttg	2580
	aaatagaagg	aatggctcaat	tgtctcccat	tctatggagt	agcagattta	aaagaatttc	2640
	taagtgtatt	atttaaacaga	aatgcacagg	atgttatgat	atgtagacct	cgaagaattga	2700
	aaagagacatt	ccaagacatt	ctctacagaa	tatccacaga	attaccocat	tacttatcta	2760
55	agtgtgttca	tgtgtgcocca	ttttttccat	tgaaagcacca	tcttcacaga	actacatgat	2820
	taaatatgtt	taagaagatt	agttaccatt	gaaattgtgt	ctgtcataaa	acagcatgag	2880
	ctcgtgtttt	aatatatctt	gtattatgtg	tcacatgtgt	atttttttaa	tgaagattca	3000
	ctgactgtgt	tttatattga	aaaaagttcc	acgtatttga	gaaaacgttaa	ataactcaat	3060
60	aac						3063

MSH2 (human) (SEQ ID NO:11)

5	MAVQPKETLQ	LESAAEVGFV	RFFQGMPEKP	TTTVRLFDRG	DFYTAHGEDA	LLAAREVFKT	60
	QGVIKYMGPA	GAKNLQSVVL	SKMNFESFVK	DLLLVRQYRV	EVYKNRAGNK	ASKENDWVLA	120
	YKASPGNLSQ	FEDILEGNND	MSASIGVVG	KMSAVDGRQ	VGVGYVDSIQ	RKLGCEPFD	180
	NDQFSNLAL	LQIGPKCEV	LPGGETAGDM	GKLRQIIQRG	GILITERKKA	DFSTKDIYQD	240
	LNRLAKKKKG	EQMGSVLP	MENQVAVSSL	SAVIKFLELL	SDSNFNGQFE	LTTFDFSGYM	300
10	KLDIAAVRAL	NLFQGSVEDT	TGSQLAALL	NKCKTPQGR	LVNQWIKQPL	MDKNRIEREL	360
	NLVEAFVEDA	ELRQTLQEDL	LRRFPDLNRL	AKKFQRQAAN	LQDCYRLYQG	INQLPNVQA	420
	LEKHEGKHQK	LLLAVFVTP	TDLRSDFSFK	QEMIETLDM	DQVENHEFLV	KPSFDPNLSE	480
	LREIMNDLEK	KMQSTLISAA	RDLGLDPGKQ	IKLDSSAQFG	YFRVTCKEE	KVLRRNNKFS	540
	TVDIQKNGVK	FTNSKLTSLN	EETKNTKEY	EEAQDAIVKE	IVNISSGVVE	PMQTLNDVLA	600
15	QLDAVVSFAH	VSNQAPVPYV	RPAILEKQGG	RILKASRHA	CVEVQDEIAF	IPNDVYFEKD	660
	KQMFHIITGP	NMGGSSTYIR	QTGVIVLMAQ	IGCFVPCESA	EVSIIVDCILA	RVGAGDSQLK	720
	GVSTFMAMEL	ETASILLRSAT	KDSLIIIDEL	GRGSTYDGF	GLAWATSEYI	ATKIGAFCMF	780
	ATHFHELTAL	ANQIPTVNNL	HVTALTTEET	LTMLYQVKKG	VCDQSGFIHV	AELANFPKHV	840
	IECAKQKALE	LEEFQYIGES	QGYDIMEPA	KKCYLEREQG	EKIIQEFLSK	VQMPFTEMS	900
20	EENITIKLQK	LKAEVIAKNN	SFVNEIISRI	KVTT			934

MSH2 (human cDNA) (SEQ ID NO:12)

	ggcgggaac	agcttagtgg	gtgtggggctc	cgccattttc	ttcaaccagg	aggtgaggag	60
25	gtctcgacat	ggcggttcgag	cogaaggaga	cgctgcagtt	ggagagcgccg	cgcgaggtcg	120
	gtctcggtcg	gtctcttcgag	ggcatgcccg	agaaagcgac	caccacagtg	cgccctttcg	180
	acggggggca	cttctatacag	cgccacggcg	aggaacggct	gctggccgcc	cgggaggtgt	240
	tcaagaccga	gggggtgagc	aagtacatgg	ggccggcagg	agcaagaagt	ctcgagagtg	300
	tgtgtcctag	taaaatgaat	tttgaatctt	ttgtaaaaag	tctctctctg	gttcgtcagt	360
	atagagttga	agtttataag	aataagagct	gaaataaagg	atccaaaggag	aatgattgtgt	420
30	atttgataat	taaggctctc	cctggcaatc	tctctcagtt	tgaagacatt	ctctcttgta	480
	acaatgatat	gtcagctctc	attggtgttg	tgggtgttaa	aatgctccga	gtgtgagccc	540
	agagacaggt	tggagtgtgg	tatgttgatt	ccatacagag	gaaactaggga	ctgtgtgaat	600
	tccttgataa	tgatcagttc	tccaatcttg	aggctctcct	catccagatt	ggaccaaaagg	660
	aattgtgtttt	accggaggga	gagactgctg	gagacatggg	gaaactgaga	caagataattc	720
35	aaagaggagg	aattctgagc	acagaaagaa	aaaaagctga	ctttcccaac	aaagacagat	780
	atcaggagcct	caaccggttg	ttgaaaggca	aaaaggggga	gcagatgaat	atgtctgtat	840
	tgccagaaga	ggagaatacag	gttgacgttt	catcactgtc	tgcggttaac	aaagtttttag	900
	aactcttatc	agatgattcc	aactttggac	agtttgaact	gactactttt	gacttcagcc	960
	agtatatgaa	attgatatatt	gcagcagtcg	gagcccttaa	ctttttctag	ggttctgttg	1020
40	aaagataccac	tggctctcag	tctctggtgt	ccttgctgaa	taagtgtaaa	accctccaag	1080
	gcagaagact	tgttaaccag	tggattaaag	agcctctcag	ggaataagaac	agaaatagag	1140
	agagattgaa	tttagtggaa	gctttgttag	aagatgcaga	attgaggcag	actttacaag	1200
	aaagatttaac	tcgtcgattc	ccagatctta	accgactctg	caagaagctt	caagaacaag	1260
	cagcaaacctt	acaagattgt	taccgactct	atcagggtat	aaatacaact	cctaattgta	1320
45	tacaggctct	ggaaaaaacat	gaaggaaaac	accgaaaatt	attgttggca	gtttttgtga	1380
	ctctcttacc	tgcattcttg	cttgactctc	ccaagtttca	ggaatagtga	gaacaaactt	1440
	tagatagtga	tcagttgtgaa	aaccatgaat	tccttgtaaa	acctctcatt	gactctaact	1500
	tcagtgaatt	aagagaataa	atgaatgact	tggaaaagaa	gatgcagtcg	atcaataata	1560
	gtgcagccag	agatcttggc	ttggaccctg	gcacacagat	taaacctggat	tcaggtgcac	1620
50	agtttggata	ttaactttcgt	gtaacctgta	aggaagaaaa	agtccttcgt	aaccaataaa	1680
	acttttagtac	tgtatagata	cagaagaatg	gtgttaaaat	taccaaacag	aaattgactt	1740
	ctttaaatga	agagataacc	aaaaataaaa	cagaatatga	agaagccagc	gatccgattg	1800
	ttaaagaaga	tgtcaatatt	tcttcaggct	atgtagaacc	ctcagagaca	ctcagatgat	1860
	tgtatagctca	gtgatagctt	gttgtcagct	gtgtcacaat	gtcaaatgga	gaacgtgttc	1920
55	catatgtgac	accagccatt	ttggagaagg	gacaagggaag	aattatatta	aaacagctca	1980
	ggcatgctgt	tgttgaagtt	caagatgaaa	ttgcatttat	tcctaagagc	gtataacttt	2040
	aaaaagatca	acagatgttc	cacatcatta	ctggcccaaaa	ctgtggaggt	aaatacaact	2100
	atattcgaca	aactgggggt	atagtactca	tggcccaaat	tggtgttttt	gtgccatgtg	2160
	agtcagcaga	agtggtccatt	gtggactgca	cttttagccg	agtaggggct	ggtgacagtc	2220

	aattgaaagg	agctctccacg	tctcatggctg	aaatgttgga	aactgcttct	atcctcaggt	2280
	ctgcaaccac	agattcattat	ataatcatag	atgaattggg	aagaggaaact	tctacctacg	2340
	atggatttgg	gttagcatgg	gctatatcag	aatacattgc	aacaaagatt	ggtgcttttt	2400
5	gcattgttgc	aaccattttt	catgaacotta	ctgccttgcc	caatcagata	ccaactgtta	2460
	ataatctaca	tgctcaagca	ctcaccactg	aagagacott	aactatgctt	tatcagggtga	2520
	agaagaagtgt	ctgtgatcaa	agtttttgga	tctatgttgc	agagcttgct	aatttcccta	2580
	agcattgta	agagctgtct	aaacagaaag	ccttggaact	tgaggagttt	cagtatatgt	2640
	gagaaatgca	aggattatgat	atcatggaac	cagcagcaaa	gaagtgtctat	ctggaaagag	2700
10	agcagaatg	aaaaattatt	caggagttcc	tgctcaagtg	gaacaaatg	ccctttactg	2760
	aaatgtcaga	agaaatacat	acaataaagt	taaaacagct	aaaagctgaa	gtaaatgcaa	2820
	agaaataagt	ctttgttaatt	gaatacattt	cacgaataaa	agttactactg	tgaataatcc	2880
	cagtaaatgga	atgaaggtaa	tattgataag	ctattgtctg	taatatgtttt	atattgtttt	2940
	atattaaccc	tttttccata	gtgttaactg	ctagtgccca	tggtgtccca	acttaataag	3000
	atatttagta	atattttact	ttgaggacat	tttcaaagat	ttttattttg	aaaaatgaga	3060
15	gctgttaactg	aggactgttt	gcaattgaca	taggcaataa	taagtgatgt	gctgaatttt	3120
	ataataaaaa	tcatgtagtt	tgtgg				3145

MLH1 (human) (SEQ ID NO:13)

20	MSFVAGVIR	LDLTVNRIA	AGEVIQFAN	AIKEMIENCL	DAKSTSIQVI	VKEGGKLQI	60
	IQDNGTIGR	EDLDIVCERF	TTSKLQSFED	LASISTYGRF	GEALASISHV	AHVTTTITKTA	120
	DGRCAYRAS	SGDKLKAPPK	PCAGNQGTQI	TVEDLFYNIA	TRRKALKNPS	EYXGKILEVV	180
	GRYSVHNAGI	SFSVKKQGET	VADVRLTPNA	TVDNIRSF	GNVSRKELIA	IGCEPKTLAF	240
	KMNGYISNAN	YSVKKCIFLL	FINHRLVEST	SLRKAIETVY	AAYLPKNTHP	FLYLSLETS	300
25	QNDVNVNHP	KHEVDFLHEE	SILERVQHH	ESKLLGSNS	RMFTQTLLP	GLAGPSGDMV	360
	KSTTLTSSS	TGSSGDKVYA	HQMVRTDSRE	OKLDAFLQPL	SKPLSSQPPA	IVDEPKTDIS	420
	SGRARQDEE	MLELPAPEAV	AAKNQSLGSD	TTKGTSEMSE	KRGFTSTQNP	KRHREDSDE	480
	MVEDDSRKE	TAACTPRRRI	INLTSVLSLQ	EEINEQGHSE	LRLEMLNHHS	FVGVNPNQAL	540
	AGQTKVLKYL	NNTLKSEELF	YQILYDFAN	FGVLRLESEPA	PLFDLAMLAL	DSPESGWTEE	600
30	DGPEKGLAEY	IVEFLKKKAE	MLADYFSL	DEEGNLGLP	LLLDNYVPEL	EGLPFILRL	660
	ATEVNWDEEK	ECFESLSKEC	AMFYSIKQY	ISEESTLSGQ	QSEVPGSIPN	SWKWTVEHIV	720
	YKALRSHLPL	PKHFTFEDGNI	LQLANLPDLY	KVFERC			780

MLH1 (human) (SEQ ID NO:14)

35	cttggctctt	ctggcgccaa	aatgtcgttc	gtggcagggg	ttattcggcg	gotggacag	60
	acagtggtga	acgcgatcgc	ggcgggggaa	gttatccagc	ggccagctaa	tgctatcaaa	120
	gagatgattg	agaactgttt	agatgcaaaa	tcacaaagta	ttcaagtgat	gtttaaaggag	180
	ggagcgctga	agttgatcca	gatccaaagc	aatggcaccg	ggatcaggaa	agaagctgt	240
	gattattgat	gtgaagtgct	cactactagt	aaactgcagt	cttttaggga	tttagccagt	300
40	atttctacct	atggctttgc	aggtgaggtc	ttggccagca	taagccatgt	ggctcatgtt	360
	actattacaa	cgaaaacacg	tgatggaag	tggtgcatac	gagcagatta	ctcagatgaa	420
	aaactgaa	ccctccctaa	accatgtgct	ggcaatcaag	ggaccagat	acagggtgag	480
	gaacctttt	acaaactatg	cagagggat	aaagctttaa	aaaataccag	tgagaataat	540
	gggaagaatt	tggaagtgt	tggcaggtat	ctcagtaaca	atgcaggcat	tgttttctca	600
45	gttaaaaaac	aaggagagac	agtagctgt	gttaggacac	taectaatgc	ctacacccgt	660
	gacaatattc	gtccatcttc	tggaaatgct	gttagtcag	aactgataga	aattggatgt	720
	gaggataaaa	ccctagccct	caaaatgaat	gggtacatat	ccaatgcaaa	ctactctgat	780
	aagaagtcca	tcttcttaact	ctctatcaac	catcgtctcg	tagaatcaac	tctccttaga	840
	aaagccatag	aaacagtgta	tgacgcctat	ttggccaaaa	acacacaccc	attctctgat	900
50	ctcagtttag	aaatcagttc	ccagaatgtg	gatgttaagt	tgacacccac	aaagcatgaa	960
	gttcactctc	tgacagagga	gagcatcctg	gagcgggtgc	agcagacat	cgagagcaag	1020
	ctcctggctg	ccaattctct	caggatgtac	ttcaccacga	cttttgtacc	aggactctgt	1080
	ggccctctga	gggagatggt	taaatccaca	acaagtctga	cctctgtctc	taactctgga	1140
	agtagctgat	aggtctatgc	ccaccagatg	gttcgtacag	attcccgaga	acagaaagct	1200
55	gtgtaatttc	tgacgcctct	gagcaaaacc	ctgtccagtc	agcccccagg	cattgtcaca	1260
	gaggaataag	cagatatttc	tagtgccag	gctaggcaga	aagatgagga	gatgcttgaa	1320
	ctccccagcc	ctgtctgaagt	ggctgccaaa	aatcagagct	tggaagggga	tacaacaaag	1380
	ggactctcag	aaatgtcaga	gaagagagga	ctactctcca	gcaacccag	aaagagacat	1440
	cgggaagatt	ctgatgtgga	aatggtggaa	gatgattccc	gaaaggaaat	gactgcagct	1500

	tgtagcttcca	ggaaga	aaatt	1560			
	aatgagcagg	gacatgaggt	tctccgggag	atgttgcata	accactccctt	cgtagggctgt	1620
	gtgaatccctc	agtgggcott	ggcacagcat	caaaccaagt	tatacctttc	caacaccacc	1680
	aaagcttagtg	aagaactgtt	ctaccagata	ctcatttatg	attttgccaa	ttttgggtgtt	1740
5	ctcagggttat	cgagggcagc	accgctcttt	gacottgcc	tgcttgccct	agatagtcca	1800
	gagagtggct	ggacagagga	agatgggtccc	aaagaaggac	tgtctgaata	catgtgttag	1860
	tttctgaaga	agaaggctga	gatgcttgca	gactatttct	ctttggaaat	tgatgaggaa	1920
	gggaacctga	ttggattacc	ccttctgatt	gacaactatg	tgcccccttt	ggaggggactg	1980
	cotattcttca	ttcttcgact	agccactgag	gtgaattggg	acgaagaaaa	ggaatgtttt	2040
10	gaagccctca	gtaaaagatg	cgctatgttc	tattccatcc	ggaagcagta	catatctgag	2100
	gactgcagccc	tctcaggcca	gcagagtga	gtgcctggct	ccattccaaa	ctctctggaag	2160
	tggaactgtg	aaacacattgt	ctataaagcc	ttgcgctcac	acattctgcc	tcctaaacat	2220
	ttcacagaag	atggaaat	at	cctgcagctt	gctaacctgc	ctgatctata	2280
	gcagaggtgtt	aaatatgtgt	atttatgcac	tgtgggatgt	gttctctttt	ctctgtattc	2340
15	cgatacaaa	tggtgtatca	aagtgtgata	tacaaagtgt	accaacataa	gtgttggtag	2400
	cacttaagac	ttatacttgc	cttctgatag	tattccttta	tacacagtgg	attgattata	2460
	aataaataga	tgtgtcttaa	cata				2484

hPMS2-134 (human) (SEQ ID NO:15)

20	MERAESSSTE	PAKAIKPIDR	KSVHQICSGQ	VVLSLSTAVK	ELVENS LDAG	ATNIDLKLDK	60
	YGVDLLEIVSD	NGCGVEENF	EGLTLKHHTS	KIQEFADLTQ	VETFGFRGEA	LSSLCALSDV	120
	TIISTCHASAK	VGT					133

hPMS2-134 (human cDNA) (SEQ ID NO:16)

25	cgaggcggat	cggtgtgtgc	atccatggag	cgagctgaga	gtcgcagtag	agaacctgct	60
	aaggccatca	aacctattga	tcggaagtca	gtccatcaga	tttgctctgg	cgaggtggta	120
	ctgagcttaa	cgactgcggt	aaaggagtta	gtagaaaaca	gtctggatgc	tggtgccact	180
	aatattgac	taaaagcttaa	ggactatgga	gtggatctta	ttgaagtgtc	agacaattga	240
	tgtgggtag	aagaagaaaa	cttcgaagcc	ttaactctga	aacatcacac	atctaagatt	300
30	caagagtttg	cgacactaac	tcaggttgaa	acttttggct	tctgggggga	agctctgagc	360
	tcactttgtg	cactgagcga	gtgcaccatt	tctacctgcc	acgcacatgc	gaaggttgga	420
	acttga						426

For further information on the background of the invention the following references

35 may be consulted, each of which is incorporated herein by reference in its entirety:

1. Glaser, V. (1996) Can ReoPro repolish tarnished monoclonal therapeutics? *Nat. Biotechnol.* 14:1216-1217.
2. Weiner, L.M. (1999) Monoclonal antibody therapy of cancer. *Semin. Oncol.* 26:43-51.
3. Saez-Llorens, X.E. *et al.* (1998) Safety and pharmacokinetics of an intramuscular humanized monoclonal antibody to respiratory syncytial virus in premature infants and infants with bronchopulmonary dysplasia. *Pediat. Infect. Dis. J.* 17:787-791.
4. Shield, C.F. *et al.* (1996) A cost-effective analysis of OKT3 induction therapy in cadaveric kidney transplantation. *Am. J. Kidney Dis.* 27:855-864.

5. Khazaeli, M.B. *et al.* (1994) Human immune response to monoclonal antibodies. *J. Immunother.* 15:42-52.
6. Emery, S.C. and W.J. Harris "Strategies for humanizing antibodies" In: ANTIBODY ENGINEERING C.A.K. Borrebaeck (Ed.) Oxford University Press, N.Y. 1995, pp. 159-183.
7. U.S. Patent No. 5,530,101 to Queen and Selick.
8. Reff, M.E. (1993) High-level production of recombinant immunoglobulins in mammalian cells. *Curr. Opin. Biotechnol.* 4:573-576.
9. Neuberger, M. and M. Gruggermann, (1997) Monoclonal antibodies. Mice perform a human repertoire. *Nature* 386:25-26.
10. Fiedler, U. and U. Conrad (1995) High-level production and long-term storage of engineered antibodies in transgenic tobacco seeds. *Bio/Technology* 13:1090-1093.
11. Baker S.M. *et al.* (1995) Male defective in the DNA mismatch repair gene PMS2 exhibit abnormal chromosome synapsis in meiosis. *Cell* 82:309-319.
15. 12. Bronner, C.E. *et al.* (1994) Mutation in the DNA mismatch repair gene homologue *hMLH1* is associated with hereditary non-polyposis colon cancer. *Nature* 368:258-261.
13. de Wind N. *et al.* (1995) Inactivation of the mouse *Msh2* gene results in mismatch repair deficiency, methylation tolerance, hyperrecombination, and predisposition to cancer. *Cell* 82:321-300.
20. 14. Drummond, J.T. *et al.* (1995) Isolation of an hMSH2-p160 heterodimer that restores mismatch repair to tumor cells. *Science* 268:1909-1912.
15. Modrich, P. (1994) Mismatch repair, genetic stability, and cancer. *Science* 266:1959-1960.
25. 16. Nicolaides, N.C. *et al.* (1998) A Naturally Occurring hPMS2 Mutation Can Confer a Dominant Negative Mutator Phenotype. *Mol. Cell. Biol.* 18:1635-1641.
17. Prolla, T.A. *et al.* (1994) MLH1, PMS1, and MSH2 Interaction during the initiation of DNA mismatch repair in yeast. *Science* 264:1091-1093.
18. Strand, M. *et al.* (1993) Destabilization of tracts of simple repetitive DNA in yeast by mutations affecting DNA mismatch repair. *Nature* 365:274-276.
- 30

19. Su, S.S., R.S. Lahue, K.G. Au, and P. Modrich (1988) Mismatch specificity of methyl directed DNA mismatch corrections in vitro. *J. Biol. Chem.* 263:6829-6835.
20. Parsons, R. *et al.* (1993) Hypermutability and mismatch repair deficiency in RER⁺ tumor cells. *Cell* 75:1227-1236.
- 5 21. Papadopoulos, N. *et al.* (1993) Mutation of a mutL homolog is associated with hereditary colon cancer. *Science* 263:1625-1629.
22. Perucho, M. (1996) Cancer of the microsatellite mutator phenotype. *Biol. Chem.* 377:675-684.
23. Nicolaides N.C., K.W. Kinzler, and B. Vogelstein (1995) Analysis of the 5' region of PMS2 reveals heterogenous transcripts and a novel overlapping gene. *Genomics* 10 29:329-334.
24. Nicolaides, N.C. *et al.* (1995) Genomic organization of the human PMS2 gene family. *Genomics* 30:195-206.
25. Palombo, F. *et al.* (1994) Mismatch repair and cancer. *Nature* 36:417.
- 15 26. Eshleman J.R. and S.D. Markowitz (1996) Mismatch repair defects in human carcinogenesis. *Hum. Mol. Genet.* 5:1489-494.
27. Liu, T. *et al.* (2000) Microsatellite instability as a predictor of a mutation in a DNA mismatch repair gene in familial colorectal cancer. *Genes Chromosomes Cancer* 27:17-25.
- 20 28. Nicolaides, N.C. *et al.* (1992) The Jun family members, c-JUN and JUND, transactivate the human *c-myc* promoter via an Ap1 like element. *J. Biol. Chem.* 267:19665-19672.
29. Shields, R.L. *et al.* (1995) Anti-IgE monoclonal antibodies that inhibit allergen-specific histamine release. *Int. Arch. Allergy Immunol.* 107:412-413.
- 25 30. Frigerio L. *et al.* (2000) Assembly, secretion, and vacuolar delivery of a hybrid immunoglobulin in plants. *Plant Physiol.* 123:1483-1494.
31. Bignami M, (2000) Unmasking a killer: DNA O(6)-methylguanine and the cytotoxicity of methylating agents. *Mutat. Res.* 462:71-82.

32. Drummond, J.T. *et al.* (1996) Cisplatin and adriamycin resistance are associated with MutLa and mismatch repair deficiency in an ovarian tumor cell line. *J. Biol. Chem.* 271:9645-19648.
33. Galio, L. *et al.* (1999) ATP hydrolysis-dependent formation of a dynamic ternary nucleoprotein complex with MutS and MutL. *Nucl. Acids Res.* 27:2325-23231.

The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific examples which are provided herein for purposes of illustration only, and are not intended to limit the scope of the invention.

EXAMPLE 1: Stable expression of dominant negative MMR genes in hybridoma cells

It has been previously shown by Nicolaides *et al.* (Nicolaides *et al.* (1998) A Naturally Occurring hPMS2 Mutation Can Confer a Dominant Negative Mutator Phenotype *Mol. Cell. Biol.* 18:1635-1641) that the expression of a dominant negative allele in an otherwise MMR proficient cell could render these host cells MMR deficient. The creation of MMR deficient cells can lead to the generation of genetic alterations throughout the entire genome of a host organisms offspring, yielding a population of genetically altered offspring or siblings that may produce biochemicals with altered properties. This patent application teaches of the use of dominant negative MMR genes in antibody-producing cells, including but not limited to rodent hybridomas, human hybridomas, chimeric rodent cells producing human immunoglobulin gene products, human cells expressing immunoglobulin genes, mammalian cells producing single chain antibodies, and prokaryotic cells producing mammalian immunoglobulin genes or chimeric immunoglobulin molecules such as those contained within single-chain antibodies. The cell expression systems described above that are used to produce antibodies are well known by those skilled in the art of antibody therapeutics.

To demonstrate the ability to create MMR defective hybridomas using dominant negative alleles of MMR genes, we first transfected a mouse hybridoma cell line that is known to produce an antibody directed against the human IgE protein with an expression vector containing the human PMS2 (cell line referred to as HBPMS2), the previously

published dominant negative PMS2 mutant referred herein as PMS134 (cell line referred to as HB134), or with no insert (cell line referred to as HBvec). The results showed that the PMS134 mutant could indeed exert a robust dominant negative effect, resulting in biochemical and genetic manifestations of MMR deficiency. Unexpectedly was the finding that the full length PMS2 also resulted in a lower MMR activity while no effect was seen in cells containing the empty vector. A brief description of the methods is provided below.

The MMR proficient mouse H36 hybridoma cell line was transfected with various *hPMS2* expression plasmids plus reporter constructs for assessing MMR activity. The MMR genes were cloned into the pEF expression vector, which contains the elongation factor promoter upstream of the cloning site followed by a mammalian polyadenylation signal. This vector also contains the NEO^r gene that allows for selection of cells retaining this plasmid. Briefly, cells were transfected with 1 µg of each vector using polyliposomes following the manufacturer's protocol (Life Technologies). Cells were then selected in 0.5 mg/ml of G418 for 10 days and G418 resistant cells were pooled together to analyze for gene expression. The pEF construct contains an intron that separates the exon 1 of the EF gene from exon 2, which is juxtaposed to the 5' end of the polylinker cloning site. This allows for a rapid reverse transcriptase polymerase chain reaction (RT-PCR) screen for cells expressing the spliced products. At day 17, 100,000 cells were isolated and their RNA extracted using the trizol method as previously described (Nicolaidis N.C., Kinzler, K.W., and Vogelstein, B. (1995) Analysis of the 5' region of PMS2 reveals heterogeneous transcripts and a novel overlapping gene. *Genomics* 29:329-334). RNAs were reverse transcribed using Superscript II (Life Technologies) and PCR amplified using a sense primer located in exon 1 of the EF gene (5'-ttt cgc aac ggg ttt gcc g-3') and an antisense primer (5'-gtt tca gag tta agc ctt cg-3') centered at nt 283 of the published human PMS2 cDNA, which will detect both the full length as well as the PMS134 gene expression. Reactions were carried out using buffers and conditions as previously described (Nicolaidis, N.C., *et al.* (1995) Genomic organization of the human PMS2 gene family. *Genomics* 30:195-206), using the following amplification parameters: 94°C for 30 sec, 52°C for 2 min, 72°C for 2 min, for 30 cycles. Reactions were analyzed on agarose gels. Figure 1 shows a representative example of PMS expression in stably transduced H36 cells.

Expression of the protein encoded by these genes were confirmed via western blot using a polyclonal antibody directed to the first 20 amino acids located in the N-terminus of the protein following the procedures previously described (data not shown) (Nicolaides *et al.* (1998) A Naturally Occurring hPMS2 Mutation Can Confer a Dominant Negative Mutator Phenotype. *Mol. Cell. Biol.* 18:1635-1641.

EXAMPLE 2: hPMS134 Causes a Defect in MMR Activity and hypermutability in hybridoma cells

A hallmark of MMR deficiency is the generation of unstable microsatellite repeats in the genome of host cells. This phenotype is referred to as microsatellite instability (MI) (Modrich, P. (1994) Mismatch repair, genetic stability, and cancer *Science* 266:1959-1960; Palombo, F., *et al.* (1994) Mismatch repair and cancer *Nature* 36:417). MI consists of deletions and/or insertions within repetitive mono-, di- and/or tri nucleotide repetitive sequences throughout the entire genome of a host cell. Extensive genetic analysis eukaryotic cells have found that the only biochemical defect that is capable of producing MI is defective MMR (Strand, M., *et al.* (1993) Destabilization of tracts of simple repetitive DNA in yeast by mutations affecting DNA mismatch repair *Nature* 365:274-276; Perucho, M. (1996) Cancer of the microsatellite mutator phenotype. *Biol Chem.* 377:675-684; Eshleman J.R., and Markowitz, S.D. (1996) Mismatch repair defects in human carcinogenesis. *Hum. Mol. Genet.* 5:1489-494). In light of this unique feature that defective MMR has on promoting MI, it is now used as a biochemical marker to survey for lack of MMR activity within host cells (Perucho, M. (1996) Cancer of the microsatellite mutator phenotype. *Biol Chem.* 377:675-684; Eshleman J.R., and Markowitz, S.D. (1996) Mismatch repair defects in human carcinogenesis. *Hum. Mol. Genet.* 5:1489-494; Liu, T., *et al.* (2000) Microsatellite instability as a predictor of a mutation in a DNA mismatch repair gene in familial colorectal cancer *Genes Chromosomes Cancer* 27:17-25).

A method used to detect MMR deficiency in eukaryotic cells is to employ a reporter gene that has a polynucleotide repeat inserted within the coding region that disrupts its reading frame due to a frame shift. In the case where MMR is defective, the reporter gene will acquire random mutations (i.e. insertions and/or deletions) within the polynucleotide

repeat yielding clones that contain a reporter with an open reading frame. We have employed the use of an MMR-sensitive reporter gene to measure for MMR activity in HBvec, HBPMS2, and HBPMS134 cells. The reporter construct used the pCAR-OF, which contains a hygromycin resistance (HYG) gene plus a β -galactosidase gene containing a 29 bp out-of-frame poly-CA tract at the 5' end of its coding region. The pCAR-OF reporter would not generate β -galactosidase activity unless a frame-restoring mutation (*i.e.*, insertion or deletion) arose following transfection. HBvec, HBPMS2, and HB134 cells were each transfected with pCAR-OF vector in duplicate reactions following the protocol described in Example 1. Cells were selected in 0.5 mg/ml G418 and 0.5mg/ml HYG to select for cells retaining both the MMR effector and the pCAR-OF reporter plasmids. All cultures transfected with the pCAR vector resulted in a similar number of HYG/G418 resistant cells. Cultures were then expanded and tested for β -galactosidase activity *in situ* as well as by biochemical analysis of cell extracts. For *in situ* analysis, 100,000 cells were harvested and fixed in 1% glutaraldehyde, washed in phosphate buffered saline solution and incubated in 1 ml of X-gal substrate solution [0.15 M NaCl, 1 mM MgCl₂, 3.3 mM K₄Fe(CN)₆, 3.3 mM K₃Fe(CN)₆, 0.2% X-Gal] in 24 well plates for 2 hours at 37°C. Reactions were stopped in 500 mM sodium bicarbonate solution and transferred to microscope slides for analysis. Three fields of 200 cells each were counted for blue (β -galactosidase positive cells) or white (β -galactosidase negative cells) to assess for MMR inactivation. Table 1 shows the results from these studies. While no β -galactosidase positive cells were observed in HBvec cells, 10% of the cells per field were β -galactosidase positive in HB134 cultures and 2% of the cells per field were β -galactosidase positive in HBPMS2 cultures.

Cell extracts were prepared from the above cultures to measure β -galactosidase using a quantitative biochemical assay as previously described (Nicolaidis *et al.* (1998)

A Naturally Occurring hPMS2 Mutation Can Confer a Dominant Negative Mutator Phenotype *Mol. Cell. Biol.* 18:1635-1641; Nicolaidis, N.C., *et al.* (1992) The Jun family members, c-JUN and JUND, transactivate the human *c-myc* promoter via an Ap1 like element. *J. Biol. Chem.* 267:19665-19672). Briefly, 100,000 cells were collected, centrifuged and resuspended in 200 μ l of 0.25M Tris, pH 8.0. Cells were

lysed by freeze/thawing three times and supernatants collected after microfugation at 14,000 rpms to remove cell debris. Protein content was determined by spectrophotometric analysis at OD^{280} . For biochemical assays, 20 μ g of protein was added to buffer containing 45 mM 2-mercaptoethanol, 1mM $MgCl_2$, 0.1 M $NaPO_4$, and 0.6 mg/ml Chlorophenol red- β -D-galactopyranoside (CPRG, Boehringer Mannheim). Reactions were incubated for 1 hour, terminated by the addition of 0.5 M Na_2CO_3 , and analyzed by spectrophotometry at 576 nm. H36 cell lysates were used to subtract out background. Figure 2 shows the β -galactosidase activity in extracts from the various cell lines. As shown, the HB134 cells produced the highest amount of β -galactosidase, while no activity was found in the HBvec cells containing the pCAR-OF. These data demonstrate the ability to generate MMR defective hybridoma cells using dominant negative MMR gene alleles.

Table 1. β -galactosidase expression of HBvec, HBPMS2 and HB134 cells transfected with pCAR-OF reporter vectors. Cells were transfected with the pCAR-OF β -galactosidase reporter plasmid. Transfected cells were selected in hygromycin and G418, expanded and stained with X-gal solution to measure for β -galactosidase activity (blue colored cells). 3 fields of 200 cells each were analyzed by microscopy. The results below represent the mean +/- standard deviation of these experiments.

Table 1.

CELL LINE	# BLUE CELLS
HBvec	0 +/- 0
HBPMS2	4 +/- 1
HB134	20 +/- 3

EXAMPLE 3: Screening strategy to identify hybridoma clones producing antibodies with higher binding affinities and/or increased immunoglobulin production.

An application of the methods presented within this document is the use of MMR deficient hybridomas or other immunoglobulin producing cells to create genetic alterations within an immunoglobulin gene that will yield antibodies with altered biochemical properties.

An illustration of this application is demonstrated within this example whereby the HB134 hybridoma (see Example 1), which is a MMR-defective cell line that produces an anti-human immunoglobulin type E (hIgE) MAb, is grown for 20 generations and clones are isolated in 96-well plates and screened for hIgE binding. Figure 3 outlines the screening procedure to identify clones that produce high affinity MAbs, which is presumed to be due to an alteration within the light or heavy chain variable region of the protein. The assay employs the use of a plate Enzyme Linked Immunosorbant Assay (ELISA) to screen for clones that produce high-affinity MAbs. 96-well plates containing single cells from HBvec or HB134 pools are grown for 9 days in growth medium (RPMI 1640 plus 10% fetal bovine serum) plus 0.5 mg/ml G418 to ensure clones retain the expression vector. After 9 days, plates are screened using an hIgE plate ELISA, whereby a 96 well plate is coated with 50µl of a 1µg/ml hIgE solution for 4 hours at 4°C. Plates are washed 3 times in calcium and magnesium free phosphate buffered saline solution (PBS⁻) and blocked in 100µl of PBS⁻ with 5% dry milk for 1 hour at room temperature. Wells are rinsed and incubated with 100 µl of a PBS solution containing a 1:5 dilution of conditioned medium from each cell clone for 2 hours. Plates are then washed 3 times with PBS⁻ and incubated for 1 hour at room temperature with 50 µl of a PBS⁻ solution containing 1:3000 dilution of a sheep anti-mouse horse radish peroxidase (HRP) conjugated

secondary antibody. Plates are then washed 3 times with PBS⁺ and incubated with 50 µls of TMB-HRP substrate (BioRad) for 15 minutes at room temperature to detect amount of antibody produced by each clone. Reactions are stopped by adding 50 µls of 500mM sodium bicarbonate and analyzed by OD at 415nm using a BioRad plate reader. Clones exhibiting an enhanced signal over background cells (H36 control cells) are then isolated and expanded into 10 ml cultures for additional characterization and confirmation of ELISA data in triplicate experiments. ELISAs are also performed on conditioned (CM) from the same clones to measure total Ig production within the conditioned medium of each well. Clones that produce an increased ELISA signal and have increased antibody levels are then further analyzed for variants that over-express and/or over-secrete antibodies as described in Example 4. Analysis of five 96-well plates each from HBvec or HB134 cells have found that a significant number of clones with a higher Optical Density (OD) value is observed in the MMR-defective HB134 cells as compared to the HBvec controls. Figure 4 shows a representative example of HB134 clones producing antibodies that bind to specific antigen (in this case IgE) with a higher affinity. Figure 4 provides raw data from the analysis of 96 wells of HBvec (left graph) or HB134 (right graph) which shows 2 clones from the HB134 plate to have a higher OD reading due to 1) genetic alteration of the antibody variable domain that leads to an increased binding to IgE antigen, or 2) genetic alteration of a cell host that leads to over-production/secretion of the antibody molecule. Anti-Ig ELISA found that the two clones, shown in figure 4 have Ig levels within their CM similar to the surrounding wells exhibiting lower OD values. These data suggest that a genetic alteration occurred within the antigen binding domain of the antibody which in turn allows for higher binding to antigen.

Clones that produced higher OD values as determined by ELISA were further analyzed at the genetic level to confirm that mutations within the light or heavy chain variable region have occurred that lead to a higher binding affinity hence yielding to a stronger ELISA signal. Briefly, 100,000 cells are harvested and extracted for RNA using the Triazol method as described above. RNAs are reverse transcribed using Superscript II as suggested by the manufacturer (Life Technology) and PCR amplified for the antigen binding sites contained within the variable light and heavy chains. Because of the heterogeneous nature of these

genes, the following degenerate primers are used to amplify light and heavy chain alleles from the parent H36 strain.

Light chain sense: 5'-GGA TTT TCA GGT GCA GAT TTT CAG-3' (SEQ ID NO:1)

Light chain antisense: 5'-ACT GGA TGG TGG GAA GAT GGA-3' (SEQ ID NO:2)

Heavy chain sense: 5'-A(G/T) GTN (A/C)AG CTN CAG (C/G)AG TC-3' (SEQ ID NO:3)

Heavy chain antisense: 5'-TNC CTT G(A/G)C CCC AGT A(G/A)(A/T)C-3' (SEQ ID NO:4)

PCR reactions using degenerate oligonucleotides are carried out at 94°C for 30 sec, 52°C for 1 min, and 72°C for 1 min for 35 cycles. Products are analyzed on agarose gels. Products of the expected molecular weights are purified from the gels by Gene Clean (Bio 101), cloned into T-tailed vectors, and sequenced to identify the wild type sequence of the variable light and heavy chains. Once the wild type sequence has been determined, non-degenerate primers were made for RT-PCR amplification of positive HB134 clones. Both the light and heavy chains were amplified, gel purified and sequenced using the corresponding sense and antisense primers. The sequencing of RT-PCR products gives representative sequence data of the endogenous immunoglobulin gene and not due to PCR induced mutations. Sequences from clones were then compared to the wild type sequence for sequence comparison. An example of the ability to create *in vivo* mutations within an immunoglobulin light or heavy chain is shown in figure 5, where HB134 clone92 was identified by ELISA to have an increased signal for hIgE. The light chain was amplified using specific sense and antisense primers. The light chain was RT-PCR amplified and the resulting product was purified and analyzed on an automated ABI377 sequencer. As shown in clone A, a residue -4 upstream of the CDR region 3 had a genetic change from ACT to TCT, which results in a Thr to Ser change within the framework region just preceding the CDR#3. In clone B, a residue -6 upstream of the CDR region had a genetic change from CCC to CTC, which results in a Pro to His change within framework region preceding CDR#2.

The ability to generate random mutations in immunoglobulin genes or chimeric immunoglobulin genes is not limited to hybridomas. Nicolaides et al. (Nicolaides *et al.* (1998) A Naturally Occurring hPMS2 Mutation Can Confer a Dominant Negative Mutator Phenotype *Mol. Cell. Biol.* 18:1635-1641) has previously shown the ability to generate hypermutable hamster cells and produce mutations within an endogenous gene. A common method for producing humanized antibodies is to graft CDR sequences from a MAbs (produced by immunizing a rodent host) onto a human Ig backbone, and transfection of the chimeric genes into Chinese Hamster Ovary (CHO) cells which in turn produce a functional Ab that is secreted by the CHO cells (Shields, R.L., *et al.* (1995) Anti-IgE monoclonal antibodies that inhibit allergen-specific histamine release. *Int. Arch. Allergy Immunol.* 107:412-413). The methods described within this application are also useful for generating genetic alterations within Ig genes or chimeric Igs transfected within host cells such as rodent cell lines, plants, yeast and prokaryotes (Frigerio L, *et al.* (2000) Assembly, secretion, and vacuolar delivery of a hybrid immunoglobulin in plants. *Plant Physiol.* 123:1483-1494).

These data demonstrate the ability to generate hypermutable hybridomas, or other Ig producing host cells that can be grown and selected, to identify structurally altered immunoglobulins yielding antibodies with enhanced biochemical properties, including but not limited to increased antigen binding affinity. Moreover, hypermutable clones that contain missense mutations within the immunoglobulin gene that result in an amino acid change or changes can be then further characterized for *in vivo* stability, antigen clearance, on-off binding to antigens, etc. Clones can also be further expanded for subsequent rounds of *in vivo* mutations and can be screened using the strategy listed above.

The use of chemical mutagens to produce genetic mutations in cells or whole organisms are limited due to the toxic effects that these agents have on "normal" cells. The use of chemical mutagens such as MNU in MMR defective organisms is much more tolerable yielding to a 10 to 100 fold increase in genetic mutation over MMR deficiency alone (Bignami M, (2000) Unmasking a killer: DNA O(6)-methylguanine and the cytotoxicity of methylating agents. *Mutat. Res.* 462:71-82). This strategy allows for the use of chemical mutagens to be used in MMR-defective Ab producing cells as a method for increasing

additional mutations within immunoglobulin genes or chimeras that may yield functional Abs with altered biochemical properties such as enhanced binding affinity to antigen, etc.

Example 4: Generation of antibody producing cells with enhanced antibody production

Analysis of clones from H36 and HB134 following the screening strategy listed above has identified a significant number of clones that produce enhanced amounts of antibody into the medium. While a subset of these clones gave higher Ig binding data as determined by ELISA as a consequence of mutations within the antigen binding domains contained in the variable regions, others were found to contain "enhanced" antibody production. A summary of the clones producing enhanced amounts of secreted MAb is shown in TABLE 2, where a significant number of clones from HB134 cells were found to produce enhanced Ab production within the conditioned medium as compared to H36 control cells.

TABLE 2. Generation of hybridoma cells producing high levels of antibody. HB134 clones were assayed by ELISA for elevated Ig levels. Analysis of 480 clones showed that a significant number of clones had elevated MAb product levels in their CM. Quantification showed that several of these clones produced greater than 500ngs/ml of MAb due to either enhanced expression and/or secretion as compared to clones from the H36 cell line.

Table 2. Production of MAb in CM from H36 and HB134 clones.

Cell Line	% clones > 400 ng/ml	% clones >500 ng/ml
H36	1/480 = 0.2%	0/480 = 0%
HB134	50/480 = 10%	8/480 = 1.7%

Cellular analysis of HB134 clones with higher MAb levels within the conditioned medium (CM) were analyzed to determine if the increased production was simply due to genetic alterations at the Ig locus that may lead to over-expression of the polypeptides forming the antibody, or due to enhanced secretion due to a genetic alteration affecting secretory pathway mechanisms. To address this issue, we expanded three HB134 clones that had increased levels of antibody within their CM. 10,000 cells were prepared for western blot

analysis to assay for intracellular steady state Ig protein levels (Figure 6). In addition, H36 cells were used as a standard reference (Lane 2) and a rodent fibroblast (Lane 1) was used as an Ig negative control. Briefly, cells were pelleted by centrifugation and lysed directly in 300 μ l of SDS lysis buffer (60 mM Tris, pH 6.8, 2% SDS, 10% glycerol, 0.1 M

5 2-mercaptoethanol, 0.001% bromophenol blue) and boiled for 5 minutes. Lysate proteins were separated by electrophoresis on 4-12% NuPAGE gels (for analysis of Ig heavy chain. Gels were electroblotted onto Immobilon-P (Millipore) in 48 mM Tris base, 40 mM glycine, 0.0375% SDS, 20% methanol and blocked at room temperature for 1 hour in Tris-buffered saline (TBS) plus 0.05% Tween-20 and 5% condensed milk. Filters were probed with a
10 1:10,000 dilution of sheep anti-mouse horseradish peroxidase conjugated monoclonal antibody in TBS buffer and detected by chemiluminescence using Supersignal substrate (Pierce). Experiments were repeated in duplicates to ensure reproducibility. Figure 6 shows a representative analysis where a subset of clones had enhanced Ig production which accounted for increased Ab production (Lane 5) while others had a similar steady state level as the
15 control sample, yet had higher levels of Ab within the CM. These data suggest a mechanism whereby a subset of HB134 clones contained a genetic alteration that in turn produces elevated secretion of antibody.

The use of chemical mutagens to produce genetic mutations in cells or whole organisms are limited due to the toxic effects that these agents have on "normal" cells. The
20 use of chemical mutagens such as MNU in MMR defective organisms is much more tolerable yielding to a 10 to 100 fold increase in genetic mutation over MMR deficiency alone (Bignami M, (2000) Unmasking a killer: DNA O(6)-methylguanine and the cytotoxicity of methylating agents. *Mutat. Res.* 462:71-82). This strategy allows for the use of chemical mutagens to be used in MMR-defective Ab producing cells as a method for increasing
25 additional mutations within immunoglobulin genes or chimeras that may yield functional Abs with altered biochemical properties such as enhanced binding affinity to antigen, etc.

Example 5: establishment of genetic stability in hybridoma cells with new output trait.

The initial steps of MMR are dependent on two protein complexes, called MutS α and
30 MutL α (Nicolaidis *et al.* (1998) A Naturally Occurring hPMS2 Mutation Can Confer a

Dominant Negative Mutator Phenotype. *Mol. Cell. Biol.* 18:1635-1641). Dominant negative MMR alleles are able to perturb the formation of these complexes with downstream biochemicals involved in the excision and polymerization of nucleotides comprising the "corrected" nucleotides. Examples from this application show the ability of a truncated MMR allele (PMS134) as well as a full length human PMS2 when expressed in a hybridoma cell line is capable of blocking MMR resulting in a hypermutable cell line that gains genetic alterations throughout its entire genome per cell division. Once a cell line is produced that contains genetic alterations within genes encoding for an antibody, a single chain antibody, over expression of immunoglobulin genes and/or enhanced secretion of antibody, it is desirable to restore the genomic integrity of the cell host. This can be achieved by the use of inducible vectors whereby dominant negative MMR genes are cloned into such vectors, introduced into Ab producing cells and the cells are cultured in the presence of inducer molecules and/or conditions. Inducible vectors include but are not limited to chemical regulated promoters such as the steroid inducible MMTV, tetracycline regulated promoters, temperature sensitive MMR gene alleles, and temperature sensitive promoters.

The results described above lead to several conclusions. First, expression of hPMS2 and PMS134 results in an increase in microsatellite instability in hybridoma cells. That this elevated microsatellite instability is due to MMR deficiency was proven by evaluation of extracts from stably transduced cells. The expression of PMS134 results in a polar defect in MMR, which was only observed using heteroduplexes designed to test repair from the 5' direction (no significant defect in repair from the 3' direction was observed in the same extracts) (Nicolaidis *et al.* (1998) A Naturally Occurring hPMS2 Mutation Can Confer a Dominant Negative Mutator Phenotype. *Mol. Cell. Biol.* 18:1635-1641). Interestingly, cells deficient in hMLH1 also have a polar defect in MMR, but in this case preferentially affecting repair from the 3' direction (Drummond, J.T., *et al.* (1996) Cisplatin and adriamycin resistance are associated with MutLa and mismatch repair deficiency in an ovarian tumor cell line. *J. Biol. Chem.* 271:9645-19648). It is known from previous studies in both prokaryotes and eukaryotes that the separate enzymatic components mediate repair from the two different directions. Our results, in combination with those of Drummond *et al.* (Shields, R.L., *et al.* (1995) Anti-IgE monoclonal antibodies that inhibit allergen-specific histamine release. *Int.*

Arch Allergy Immunol. 107:412-413), strongly suggest a model in which 5' repair is primarily dependent on hPMS2 while 3' repair is primarily dependent on hMLH1. It is easy to envision how the dimeric complex between PMS2 and MLH1 might set up this directionality. The combined results also demonstrate that a defect in directional MMR is sufficient to produce a MMR defective phenotype and suggests that any MMR gene allele is useful to produce genetically altered hybridoma cells, or a cell line that is producing Ig gene products. Moreover, the use of such MMR alleles will be useful for generating genetically altered Ig polypeptides with altered biochemical properties as well as cell hosts that produce enhanced amounts of antibody molecules.

Another method that is taught in this application is that ANY method used to block MMR can be performed to generate hypermutability in an antibody-producing cell that can lead to genetically altered antibodies with enhanced biochemical features such as but not limited to increased antigen binding, enhanced pharmacokinetic profiles, etc. These processes can also be used to generate antibody producer cells that have increased Ig expression as shown in Example 4, figure 6 and/or increased antibody secretion as shown in Table 2.

In addition, we demonstrate the utility of blocking MMR in antibody producing cells to increase genetic alterations within Ig genes that may lead to altered biochemical features such as, but not limited to, increased antigen binding affinities (Figure 5A and 5B). The blockade of MMR in such cells can be through the use of dominant negative MMR gene alleles from any species including bacteria, yeast, protozoa, insects, rodents, primates, mammalian cells, and man. Blockade of MMR can also be generated through the use of antisense RNA or deoxynucleotides directed to any of the genes involved in the MMR biochemical pathway. Blockade of MMR can be through the use of polypeptides that interfere with subunits of the MMR complex including but not limited to antibodies. Finally, the blockade of MMR may be through the use chemicals such as but not limited to nonhydrolyzable ATP analogs, which have been shown to block MMR (Galio, L, *et al.* (1999) ATP hydrolysis-dependent formation of a dynamic ternary nucleoprotein complex with MutS and MutL. *Nucl. Acids Res.* 27:2325-23231).

WE CLAIM:

1. A method for making a hypermutable, antibody producing cell, comprising introducing into a cell capable of producing antibodies a polynucleotide comprising a dominant negative allele of a mismatch repair gene.
2. The method of claim 1 wherein said polynucleotide is introduced by transfection of a suspension of cells *in vitro*.
3. The method of claim 1 wherein said mismatch repair gene is *PMS2*.
4. The method of claim 1 wherein said mismatch repair gene is human *PMS2*.
5. The method of claim 1 wherein said mismatch repair gene is *MLH1*.
6. The method of claim 1 wherein said mismatch repair gene is *PMS1*.
7. The method of claim 1 wherein said mismatch repair gene is *MSH2*.
8. The method of claim 1 wherein said mismatch repair gene is *MSH1*.
9. The method of claim 4 wherein said allele comprises a truncation mutation.
10. The method of claim 4 wherein said allele comprises a truncation mutation at codon 134.
11. The method of claim 10 wherein said truncation mutation is a thymidine at nucleotide 424 of wild-type *PMS2*.
12. The method of claim 1 wherein said polynucleotide is introduced into a fertilized egg of an animal.
13. The method of claim 12 wherein the fertilized egg is subsequently implanted into a pseudo-pregnant female whereby the fertilized egg develops into a mature transgenic animal.
14. The method of claim 12 wherein said mismatch repair gene is *PMS2*.
15. The method of claim 12 wherein said mismatch repair gene is human *PMS2*.
16. The method of claim 12 wherein said mismatch repair gene is human *MLH1*.
17. The method of claim 12 wherein said mismatch repair gene is human *PMS1*.
18. The method of claim 11 wherein said mismatch repair gene is a human *mutL* homolog.
19. The method of claim 15 wherein said allele comprises a truncation mutation.
20. The method of claim 15 wherein said allele comprises a truncation mutation at codon 134.

21. The method of claim 19 wherein said truncation mutation is a thymidine at nucleotide 424 of wild-type *PMS2*.
22. The method of claim 1 wherein said capability is due to the co-introduction of an immunoglobulin gene into said cell.
23. A homogeneous culture of hypermutable, mammalian cells wherein said cells comprise a dominant negative allele of a mismatch repair gene.
24. The culture of hypermutable, mammalian cells of claim 23 wherein the mismatch repair gene is *PMS2*.
25. The culture of hypermutable, mammalian cells of claim 24 wherein the mismatch repair gene is human *PMS2*.
26. The culture of hypermutable, mammalian cells of claim 23 wherein the mismatch repair gene is *MLH1*.
27. The culture of hypermutable, mammalian cells of claim 23 wherein the mismatch repair gene is *PMS1*.
28. The culture of hypermutable, mammalian cells of claim 23 wherein the mismatch repair gene is a human *mutL* homolog.
29. The culture of hypermutable, mammalian cells of claim 23 wherein the cells express a protein consisting of the first 133 amino acids of hPMS2.
30. A method for generating a mutation in a gene affecting antibody production in an antibody-producing cell comprising:
 - growing a said cell comprising said gene and a dominant negative allele of a mismatch repair gene; and
 - testing the cell to determine whether said gene of interest harbors a mutation.
31. The method of claim 30 wherein the step of testing comprises analyzing a nucleotide sequence of said gene.
32. The method of claim 30 wherein the step of testing comprises analyzing mRNA transcribed from said gene.
33. The method of claim 30 wherein the step of testing comprises analyzing a protein encoded by the gene of interest.

34. The method of claim 30 wherein the step of testing comprises analyzing the phenotype of said gene.
35. The method of claim 30 wherein the step of testing comprises analyzing the binding activity of an antibody.
36. A method wherein a mammalian cell is made MMR defective by the process of introducing a polynucleotide comprising an antisense oligonucleotide targeted against an allele of a mismatch repair gene into a mammalian cell, whereby the cell becomes hypermutable.
37. The method of claim 36 wherein the step of testing comprises analyzing a nucleotide sequence of said gene.
38. The method of claim 36 wherein the step of testing comprises analyzing mRNA transcribed from said gene.
39. The method of claim 36 wherein the step of testing comprises analyzing a protein encoded by said gene.
40. The method of claim 36 wherein the step of testing comprises analyzing the phenotype of said gene.
41. The method of claim 36 wherein the step of testing comprises analyzing the binding activity of an antibody.
42. A method for generating a mutation in a gene affecting antibody production in an antibody-producing cell comprising:
growing said cell comprising said gene and a polynucleotide encoding a dominant negative allele of a mismatch repair gene; and
testing said cell to determine whether said cell harbors at least one mutation in said gene yielding to a new biochemical feature to the product of said gene, wherein said new biochemical feature is selected from the group consisting of over-expression of said product, enhanced secretion of said product, enhanced affinity of said product for antigen, and combinations thereof.
43. The method of claim 42 wherein the step of testing comprises analyzing the steady state expression of the immunoglobulin gene of said cell.

44. The method of claim 42 wherein the step of testing comprises analyzing steady state mRNA transcribed from the immunoglobulin gene of said cell.
45. The method of claim 42 wherein the step of testing comprises analyzing the amount of secreted protein encoded by the immunoglobulin gene of said cell.
46. The method of claim 36 wherein the cell is made by the process of introducing a polynucleotide comprising a dominant negative allele of a mismatch repair gene into a cell in the presence of DNA mutagens.
47. The method of claim 46 wherein the step of testing comprises analyzing a nucleotide sequence of an immunoglobulin gene of said cell.
48. The method of claim 46 wherein the step of testing comprises analyzing mRNA transcribed from the immunoglobulin gene of said cell.
49. The method of claim 46 wherein the step of testing comprises analyzing the immunoglobulin protein encoded by said gene.
50. The method of claim 46 wherein the step of testing comprises analyzing the biochemical activity of the protein encoded by said gene.
51. A hypermutable transgenic mammalian cell made by the method of claim 42.
52. The transgenic mammalian cell of claim 51 wherein said cell is from primate.
53. The transgenic mammalian cell of claim 51 wherein said cell is from rodent.
54. The transgenic mammalian cell of claim 51 wherein said cell is from human.
55. The transgenic mammalian cell of claim 51 wherein said cell is eukaryotic.
56. The transgenic mammalian cell of claim 51 wherein said cell is prokaryotic.
57. A method of reversibly altering the hypermutability of an antibody producing cell comprising introducing an inducible vector into a cell, wherein said inducible vector comprises a dominant negative allele of a mismatch repair gene operably linked to an inducible promoter, and inducing said cell to express said dominant negative mismatch repair gene.
58. The method of claim 57 wherein said mismatch repair gene is *PMS2*.
59. The method of claim 58 wherein said mismatch repair gene is human *PMS2*.
60. The method of claim 57 wherein said mismatch repair gene is *MLH1*.
61. The method of claim 57 wherein said mismatch repair gene is *PMS1*.

62. The method of claim 57 wherein said mismatch repair gene is a human *mutL* homolog.
63. The method of claim 57 wherein said cell expresses a protein consisting of the first 133 amino acids of hPMS2.
64. The method of claim 57 further comprising analyzing the immunoglobulin protein expressed by said antibody producing cell.
65. The method of claim 64 further comprising ceasing induction of said cell, thereby restoring genetic stability of said cell.
66. A method of producing genetically altered antibodies comprising
transfecting a polynucleotide encoding an immunoglobulin protein into a cell,
wherein said cell comprises a dominant negative mismatch repair gene;
growing said cell, thereby producing a hypermutated polynucleotide encoding
a hypermutated immunoglobulin protein;
screening for a desirable property of said hypermutated immunoglobulin
protein;
isolating said hypermutated polynucleotide; and
transfecting said hypermutated polynucleotide into a genetically stable cell,
thereby producing a hypermutated antibody-producing, genetically stable cell.
67. The method of claim 66 wherein said mismatch repair gene is *PMS2*.
68. The method of claim 66 wherein said mismatch repair gene is human *PMS2*.
69. The method of claim 66 wherein said mismatch repair gene is *MLH1*.
70. The method of claim 66 wherein said mismatch repair gene is *PMS1*.
71. The method of claim 66 wherein said mismatch repair gene is a human *mutL* homolog.
72. The method of claim 66 wherein said cell expresses a protein consisting of the first 133 amino acids of hPMS2.

ABSTRACT

Dominant negative alleles of human mismatch repair genes can be used to generate hypermutable cells and organisms. By introducing these genes into cells and transgenic animals, new cell lines and animal varieties with novel and useful properties can be prepared more efficiently than by relying on the natural rate of mutation. These methods are useful for generating genetic diversity within immunoglobulin genes directed against an antigen of interest to produce altered antibodies with enhanced biochemical activity. Moreover, these methods are useful for generating antibody-producing cells with increased level of antibody production.




FIG. 1/6




Fig. 2/6

Fig. 3/6






Fig.4/6

HBvec clones



HB134 clones



* = clones with a significant difference in antigen binding




Fig. 5A/6




Fig. 5B/6

33707468..3.407468




Fig. 6/6

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In Re Application of:

Nicholas C. Nicolaides, Luigi Grasso, and Philip
M. Sass

Group Art Unit: Not assigned**Examiner:** Not assigned

For: METHODS FOR GENERATING
GENETICALLY ALTERED ANTIBODY-
PRODUCING CELL LINES WITH
IMPROVED ANTIBODY
CHARACTERISTICS

DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name; and

I believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a



Utility Patent



Design Patent

is sought on the invention, whose title appears above, the specification of which:



is attached hereto.



was filed on _____ as Serial No. _____.



said application having been amended on _____.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the U.S. Patent and Trademark Office all information known to be material to the patentability of this application in accordance with 37 CFR § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. § 119(a-d) of any **foreign application(s)** for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of any application on which priority is claimed:

Priority Claimed (If X'd)	Country	Serial Number	Date Filed
<input type="checkbox"/>	_____	_____	_____
<input type="checkbox"/>	_____	_____	_____
<input type="checkbox"/>	_____	_____	_____
<input type="checkbox"/>	_____	_____	_____

I hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose to the U.S. Patent and Trademark Office all information known to be material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

Serial Number	Date Filed	Patented/Pending/Abandoned
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____

I hereby claim the benefit under 35 U.S.C. § 119(e) of any United States provisional application(s) listed below:

Serial Number	Date Filed
_____	_____

I hereby appoint the following persons of the firm of **WOODCOCK WASHBURN KURTZ MACKIEWICZ & NORRIS LLP**, One Liberty Place - 46th Floor, Philadelphia, Pennsylvania 19103 as attorney(s) and/or agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

John W. CaldwellReg. No. 28,937Patrick J. FarleyReg. No. 42,524

Address all telephone calls and correspondence to:

Patrick J. Farley
**WOODCOCK WASHBURN KURTZ
MACKIEWICZ & NORRIS LLP**
One Liberty Place - 46th Floor
Philadelphia PA 19103
Telephone No.: (215) 568-3100
Facsimile No.: (215) 568-3439

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name: Nicholas C. Nicolaides	
Mailing Address: 4 Cider Mill Court Boothwyn, PA 19061	Signature _____ Date of Signature: _____
City/State of Actual Residence: Boothwyn, Pennsylvania	Citizenship: <u>United States</u>

Name: Luigi Grasso	
Mailing Address: 834 Chestnut Street, Apt#816 Philadelphia, PA 19107	Signature Date of Signature: _____
City/State of Actual Residence: Philadelphia, Pennsylvania	Citizenship: <u>United States</u>

Name: Philip M. Sass	
Mailing Address: 1903 Blackhawk Circle Audubon, PA 19403	Signature Date of Signature: _____
City/State of Actual Residence: Audubon, Pennsylvania	Citizenship: <u>United States</u>

Name:	
Mailing Address:	Signature Date of Signature: _____
City/State of Actual Residence:	Citizenship: _____

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In Re Application of:

Nicholas C. Nicolaides, Luigi Grasso, and
Philip M. Sass

Serial No.: Not assigned**Group Art Unit:** Not assigned**Filing Date:** November 7, 2000**Examiner:** Not assigned

For: METHODS FOR GENERATING GENETICALLY ALTERED ANTIBODY-
PRODUCING CELL LINES WITH IMPROVED ANTIBODY
CHARACTERISTICS

BOX SEQUENCE


Assistant Commissioner for Patents
Washington DC 20231

**STATEMENT TO SUPPORT FILING AND SUBMISSION IN ACCORDANCE
WITH 37 CFR §§ 1.821 THROUGH 1.825**

- ☒ I hereby state, in accordance with the requirements of 37 C.F.R. §1.821(f), that the contents of the paper and computer readable copies of the Sequence Listing, submitted in accordance with 37 CFR §1.821(c) and (e), respectively are the same.
- ☐ I hereby state that the submission filed in accordance with 37 CFR §1.821(g) does not include new matter.
- ☐ I hereby state that the submission filed in accordance with 37 CFR §1.821(h) does not include new matter or go beyond the disclosure in the international application as filed.
- ☐ I hereby state that the amendments, made in accordance with 37 CFR §1.825(a), included in the substitute sheet(s) of the Sequence Listing are supported in the application, as filed, at pages _____. I hereby state that the substitute sheet(s) of the Sequence Listing does (do) not include new matter.
- ☐ I hereby state that the substitute copy of the computer readable form, submitted in accordance with 37 CFR §1.825(b), is the same as the amended Sequence Listing.

☐ I hereby state that the substitute copy of the computer readable form, submitted in accordance with 37 CFR §1.825(d), contains identical data to that originally filed.

Date: 11/7/00


Patrick J. Farley
Registration No. 42,524

Woodcock Washburn Kurtz
Mackiewicz & Norris LLP
One Liberty Place - 46th Floor
Philadelphia PA 19103
Telephone: (215) 568-3100
Facsimile: (215) 568-3439

© 1997 WVKMN

SEQUENCE LISTING

<110> Nicolaides, Nicholas C
Grasso, Luigi
Sass, Philip M

<120> METHODS FOR GENERATING GENETICALLY ALTERED
ANTIBODY-PRODUCING CELL LINES WITH IMPROVED ANTIBODY
CHARACTERISTICS

<130> MOR-0003

<140> 00/000,000

<141> 2000-11-07

<160> 16

<170> PatentIn Ver. 2.1

<210> 1

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:oligonucleotide
primer

<400> 1

ggattttcag gtgcagattt tcag

24

<210> 2

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:oligonucleotide
primer

<400> 2

actggatggt gggaagatgg a

21

<210> 3

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:oligonucleotide
primer

<400> 3

akgtnmagct ncagsagtc

19

<210> 4

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:oligonucleotide
primer

<400> 4

tncottgrcc ccagtarwc

19

<210> 5

<211> 859

<212> PRT

<213> Mus musculus

<400> 5

Met Glu Gln Thr Glu Gly Val Ser Thr Glu Cys Ala Lys Ala Ile Lys
1 5 10 15

Pro Ile Asp Gly Lys Ser Val His Gln Ile Cys Ser Gly Gln Val Ile
20 25 30

Leu Ser Leu Ser Thr Ala Val Lys Glu Leu Ile Glu Asn Ser Val Asp
35 40 45

Ala Gly Ala Thr Thr Ile Asp Leu Arg Leu Lys Asp Tyr Gly Val Asp
50 55 60

Leu Ile Glu Val Ser Asp Asn Gly Cys Gly Val Glu Glu Glu Asn Phe
65 70 75 80

Glu Gly Leu Ala Leu Lys His His Thr Ser Lys Ile Gln Glu Phe Ala
85 90 95

Asp Leu Thr Gln Val Glu Thr Phe Gly Phe Arg Gly Glu Ala Leu Ser

100	105	110
Ser Leu Cys Ala Leu Ser Asp Val Thr Ile Ser Thr Cys His Gly Ser 115	120	125
Ala Ser Val Gly Thr Arg Leu Val Phe Asp His Asn Gly Lys Ile Thr 130	135	140
Gln Lys Thr Pro Tyr Pro Arg Pro Lys Gly Thr Thr Val Ser Val Gln 145	150	155 160
His Leu Phe Tyr Thr Leu Pro Val Arg Tyr Lys Glu Phe Gln Arg Asn 165	170	175
Ile Lys Lys Glu Tyr Ser Lys Met Val Gln Val Leu Gln Ala Tyr Cys 180	185	190
Ile Ile Ser Ala Gly Val Arg Val Ser Cys Thr Asn Gln Leu Gly Gln 195	200	205
Gly Lys Arg His Ala Val Val Cys Thr Ser Gly Thr Ser Gly Met Lys 210	215	220
Glu Asn Ile Gly Ser Val Phe Gly Gln Lys Gln Leu Gln Ser Leu Ile 225	230	235 240
Pro Phe Val Gln Leu Pro Pro Ser Asp Ala Val Cys Glu Glu Tyr Gly 245	250	255
Leu Ser Thr Ser Gly Arg His Lys Thr Phe Ser Thr Phe Arg Ala Ser 260	265	270
Phe His Ser Ala Arg Thr Ala Pro Gly Gly Val Gln Gln Thr Gly Ser 275	280	285
Phe Ser Ser Ser Ile Arg Gly Pro Val Thr Gln Gln Arg Ser Leu Ser 290	295	300
Leu Ser Met Arg Phe Tyr His Met Tyr Asn Arg His Gln Tyr Pro Phe 305	310	315 320
Val Val Leu Asn Val Ser Val Asp Ser Glu Cys Val Asp Ile Asn Val 325	330	335
Thr Pro Asp Lys Arg Gln Ile Leu Leu Gln Glu Glu Lys Leu Leu Leu 340	345	350
Ala Val Leu Lys Thr Ser Leu Ile Gly Met Phe Asp Ser Asp Ala Asn		

355

360

365

Lys Leu Asn Val Asn Gln Gln Pro Leu Leu Asp Val Glu Gly Asn Leu
370 375 380

Val Lys Leu His Thr Ala Glu Leu Glu Lys Pro Val Pro Gly Lys Gln
385 390 395 400

Asp Asn Ser Pro Ser Leu Lys Ser Thr Ala Asp Glu Lys Arg Val Ala
405 410 415

Ser Ile Ser Arg Leu Arg Glu Ala Phe Ser Leu His Pro Thr Lys Glu
420 425 430

Ile Lys Ser Arg Gly Pro Glu Thr Ala Glu Leu Thr Arg Ser Phe Pro
435 440 445

Ser Glu Lys Arg Gly Val Leu Ser Ser Tyr Pro Ser Asp Val Ile Ser
450 455 460

Tyr Arg Gly Leu Arg Gly Ser Gln Asp Lys Leu Val Ser Pro Thr Asp
465 470 475 480

Ser Pro Gly Asp Cys Met Asp Arg Glu Lys Ile Glu Lys Asp Ser Gly
485 490 495

Leu Ser Ser Thr Ser Ala Gly Ser Glu Glu Glu Phe Ser Thr Pro Glu
500 505 510

Val Ala Ser Ser Phe Ser Ser Asp Tyr Asn Val Ser Ser Leu Glu Asp
515 520 525

Arg Pro Ser Gln Glu Thr Ile Asn Cys Gly Asp Leu Asp Cys Arg Pro
530 535 540

Pro Gly Thr Gly Gln Ser Leu Lys Pro Glu Asp His Gly Tyr Gln Cys
545 550 555 560

Lys Ala Leu Pro Leu Ala Arg Leu Ser Pro Thr Asn Ala Lys Arg Phe
565 570 575

Lys Thr Glu Glu Arg Pro Ser Asn Val Asn Ile Ser Gln Arg Leu Pro
580 585 590

Gly Pro Gln Ser Thr Ser Ala Ala Glu Val Asp Val Ala Ile Lys Met
595 600 605

Asn Lys Arg Ile Val Leu Leu Glu Phe Ser Leu Ser Ser Leu Ala Lys

610	615	620
Arg Met Lys Gln Leu	Gln His Leu Lys Ala	Gln Asn Lys His Glu Leu
625	630	635 640
Ser Tyr Arg Lys Phe	Arg Ala Lys Ile Cys	Pro Gly Glu Asn Gln Ala
645	650	655
Ala Glu Asp Glu Leu	Arg Lys Glu Ile Ser	Lys Ser Met Phe Ala Glu
660	665	670
Met Glu Ile Leu Gly	Gln Phe Asn Leu Gly	Phe Ile Val Thr Lys Leu
675	680	685
Lys Glu Asp Leu Phe	Leu Val Asp Gln His	Ala Ala Asp Glu Lys Tyr
690	695	700
Asn Phe Glu Met Leu	Gln Gln His Thr Val	Leu Gln Ala Gln Arg Leu
705	710	715 720
Ile Thr Pro Gln Thr	Leu Asn Leu Thr Ala	Val Asn Glu Ala Val Leu
725	730	735
Ile Glu Asn Leu Glu	Ile Phe Arg Lys Asn	Gly Phe Asp Phe Val Ile
740	745	750
Asp Glu Asp Ala Pro	Val Thr Glu Arg Ala	Lys Leu Ile Ser Leu Pro
755	760	765
Thr Ser Lys Asn Trp	Thr Phe Gly Pro Gln	Asp Ile Asp Glu Leu Ile
770	775	780
Phe Met Leu Ser Asp	Ser Pro Gly Val Met	Cys Arg Pro Ser Arg Val
785	790	795 800
Arg Gln Met Phe Ala	Ser Arg Ala Cys Arg	Lys Ser Val Met Ile Gly
805	810	815
Thr Ala Leu Asn Ala	Ser Glu Met Lys Lys	Leu Ile Thr His Met Gly
820	825	830
Glu Met Asp His Pro	Trp Asn Cys Pro His	Gly Arg Pro Thr Met Arg
835	840	845
His Val Ala Asn Leu	Asp Val Ile Ser Gln	Asn
850	855	

<210> 6
 <211> 3056
 <212> DNA
 <213> Mus musculus

<400> 6

gaattccggt	gaaggctcgt	aagaatttcc	agattccctga	gtatcattgg	aggagacaga	60
taacctgtcg	tcaggtaaacg	atgggtgtata	tgcaacagaa	atgggtgttcc	ctggagacgc	120
gtcttttccc	gagagcgcca	ccgcaactct	cccgcggtga	ctgtgactgg	aggagctcgt	180
catccatgga	gcaaacccgaa	ggcgtgagta	cagaatgtgc	taaggccatc	aagcctattg	240
atgggaagtc	agtcacatcaa	atttgttctg	ggcaggtgat	actcagttta	agcaccgctg	300
tgaaggagtt	gatagaaaat	agtgtagatg	ctggtgctac	tactattgat	ctaaggctta	360
aagactatgg	ggtggacctc	attgaagttt	cagacaatgg	atgtggggta	gaagaagaaa	420
actttgaagg	tctagctctg	aaacatcaca	catctaagat	tcaagagttt	gccgacctca	480
cgcagggtga	aactttcggc	tttcgggggg	aagctctgag	ctctctgtgt	gcactaagtg	540
atgtcactat	atctacctgc	cacgggtctg	caagcgttgg	gactcgactg	gtgtttgacc	600
ataatgggaa	aatcaccccg	aaaactccct	accccgacc	taaaaggaa	acagtcagtg	660
tgcagcactt	attttatata	ctacccctgc	gttacaaga	gtttcagagg	aacattaaaa	720
aggagatttc	caaaatgggt	caggctctac	aggcgtactg	tatcatctca	gcaggcgctcc	780
gtgttaagctg	cactaatcgt	ctcggacagg	ggaagcgcca	cgctgtgggt	tgccaaagcg	840
gcacgtctgg	catgaaggaa	aatatcgggt	ctgtgtttgg	ccagaagcag	ttgcaaagcc	900
tcattctctt	tggtcagctg	ccccctagtg	acgctgtgtg	tgaagagatc	ggcctgagca	960
cttcaggcag	ccacaacccc	ttttctacgt	ttcgggcttc	atttcacagt	gcagcgacgg	1020
cgcggggagg	agtgcaacag	acaggcagtt	tttcttcac	aatcagagcg	cctgtgacct	1080
agcaaaaggtc	tctaagcttg	tcaatgaggt	tttatccat	gtataaccgg	catcagtaacc	1140
cattttgtcgt	ccctaacgtt	tccgttgact	cagaatgtgt	ggatattaat	gactccagc	1200
ataaaaaggca	aattctacta	caagaagaga	agctattgct	ggcggtttta	aagacctcct	1260
tgataggaa	gtttgacagt	gatgcaacaa	agcttaattg	caaccagcag	ccactgtcag	1320
atgttgaaagg	taacttagta	aagctgcata	ctgcagaact	agaaaaacct	gtgccaggaa	1380
agcaagataa	ctctccttca	ctgaagagca	cagcagacga	gaaaagggtg	gcattccatc	1440
ccaggctagc	agagggcctt	tctcttcac	ctactaaaag	gatcaagtct	aggggtccag	1500
agactgctga	actgacacgg	agttttccaa	gtgagaaaa	ggcgctgtta	tcctcttacc	1560
cttcagacgt	catctcttac	agaggctccc	gtggctcgca	ggacaaattg	gtgagtccca	1620
cggacagccc	tggtgactgt	atggacagag	agaaaataga	aaagagctca	gggctcagca	1680
gcacctcagc	tggtctctgag	gaagagttca	gcacccca	agtgccaggt	agcttttagc	1740
gtgactataa	cgtgagctcc	ctagaagaca	gaccttctca	ggaaaaccata	aactgtgggt	1800
acctggactg	ccgtctctca	ggtacagagc	agtccttgaa	gccagaagac	catggatata	1860
aatgcaaaagc	tctaccttca	gctogtctgt	caccacacaa	tgccaagcgc	ttcaagagcag	1920
aggaaagacc	ctcaaatgtc	aacattttct	aaagattgac	tggtctctcag	agcacctcag	1980
cagctgaggt	cgatgtagcc	ataaaaaatga	ataagagaat	cgtgtctctc	gagttctctc	2040
tgagttctct	agctaagcga	atgaagcagt	tacagcacct	aaaggcgcag	aaacaacatg	2100
aactgaggtta	cagaaaaatt	agggccaa	tttgccctg	agaaaaacca	gcagcagaag	2160
atgaactcag	aaaaagagatt	agtaaatcga	tgtttgacga	gatggagatc	tttggttcagt	2220
tttaacctggg	atttatagta	accacaaactga	aagaggacct	cttctctggtg	gaccagcatg	2280
ctgcggatga	gaagtacaac	tttgagatgc	tgacgacgca	cacgggtctc	caggcgacga	2340
ggctcatcac	accccgagact	ctgaacttaa	ctgctgtcaa	tgaagctgta	ctgatagaaa	2400
atctggaaat	attcagaaaag	aatggccttg	actttgtcat	tgatgaggat	gctccagtca	2460
ctgaaagggc	taaattgatt	tccttaccac	ctagtaaaaa	ctggaccttt	ggaccccaag	2520

atatagatga actgatcttt atgttaagt acagccctgg ggtcatgtgc cggccctcac 2580
 gagtcagaca gatgtttgct tccagagcct gtcggaagtc agtgaatgatt ggaacggcgc 2640
 tcaatgcgag cgagatgaag aagctcatca cccacatggg tgagatggac cacccttgga 2700
 actgccccca cggcaggcca accatgaggc acgttgccaa tctggatgtc atctctcaga 2760
 actgacacac cccttgtagc atagagttaa ttacagattg ttcggtttgc aaagagaagg 2820
 ttttaagtaa tctgattatc gttgtacaaa aattagcatg ctgctttaat gtactggatc 2880
 catttaaaag cagtgttaag gcaggcatga tggagtgttc ctctagctca gctacttggg 2940
 tgatccggtg ggagctcatg tgagcccagg actttgagac cactccgagc cacattcatg 3000
 agactcaatt caaggacaaa aaaaaaaga tatttttgaa gccttttaaa aaaaaa 3056

<210> 7

<211> 862

<212> PRT

<213> Homo sapiens

<400> 7

Met Glu Arg Ala Glu Ser Ser Ser Thr Glu Pro Ala Lys Ala Ile Lys
 1 5 10 15

Pro Ile Asp Arg Lys Ser Val His Gln Ile Cys Ser Gly Gln Val Val
 20 25 30

Leu Ser Leu Ser Thr Ala Val Lys Glu Leu Val Glu Asn Ser Leu Asp
 35 40 45

Ala Gly Ala Thr Asn Ile Asp Leu Lys Leu Lys Asp Tyr Gly Val Asp
 50 55 60

Leu Ile Glu Val Ser Asp Asn Gly Cys Gly Val Glu Glu Glu Asn Phe
 65 70 75 80

Glu Gly Leu Thr Leu Lys His His Thr Ser Lys Ile Gln Glu Phe Ala
 85 90 95

Asp Leu Thr Gln Val Glu Thr Phe Gly Phe Arg Gly Glu Ala Leu Ser
 100 105 110

Ser Leu Cys Ala Leu Ser Asp Val Thr Ile Ser Thr Cys His Ala Ser
 115 120 125

Ala Lys Val Gly Thr Arg Leu Met Phe Asp His Asn Gly Lys Ile Ile
 130 135 140

Gln Lys Thr Pro Tyr Pro Arg Pro Arg Gly Thr Thr Val Ser Val Gln
 145 150 155 160

Gln Leu Phe Ser Thr Leu Pro Val Arg His Lys Glu Phe Gln Arg Asn

165					170					175					
Ile	Lys	Lys	Glu	Tyr	Ala	Lys	Met	Val	Gln	Val	Leu	His	Ala	Tyr	Cys
			180					185					190		
Ile	Ile	Ser	Ala	Gly	Ile	Arg	Val	Ser	Cys	Thr	Asn	Gln	Leu	Gly	Gln
		195					200					205			
Gly	Lys	Arg	Gln	Pro	Val	Val	Cys	Thr	Gly	Gly	Ser	Pro	Ser	Ile	Lys
		210					215					220			
Glu	Asn	Ile	Gly	Ser	Val	Phe	Gly	Gln	Lys	Gln	Leu	Gln	Ser	Leu	Ile
		225					230					235			240
Pro	Phe	Val	Gln	Leu	Pro	Pro	Ser	Asp	Ser	Val	Cys	Glu	Glu	Tyr	Gly
			245					250						255	
Leu	Ser	Cys	Ser	Asp	Ala	Leu	His	Asn	Leu	Phe	Tyr	Ile	Ser	Gly	Phe
			260					265					270		
Ile	Ser	Gln	Cys	Thr	His	Gly	Val	Gly	Arg	Ser	Ser	Thr	Asp	Arg	Gln
		275					280						285		
Phe	Phe	Phe	Ile	Asn	Arg	Arg	Pro	Cys	Asp	Pro	Ala	Lys	Val	Cys	Arg
		290					295					300			
Leu	Val	Asn	Glu	Val	Tyr	His	Met	Tyr	Asn	Arg	His	Gln	Tyr	Pro	Phe
		305					310					315			320
Val	Val	Leu	Asn	Ile	Ser	Val	Asp	Ser	Glu	Cys	Val	Asp	Ile	Asn	Val
			325						330				335		
Thr	Pro	Asp	Lys	Arg	Gln	Ile	Leu	Leu	Gln	Glu	Glu	Lys	Leu	Leu	Leu
			340					345					350		
Ala	Val	Leu	Lys	Thr	Ser	Leu	Ile	Gly	Met	Phe	Asp	Ser	Asp	Val	Asn
		355					360						365		
Lys	Leu	Asn	Val	Ser	Gln	Gln	Pro	Leu	Leu	Asp	Val	Glu	Gly	Asn	Leu
		370					375					380			
Ile	Lys	Met	His	Ala	Ala	Asp	Leu	Glu	Lys	Pro	Met	Val	Glu	Lys	Gln
		385					390					395			400
Asp	Gln	Ser	Pro	Ser	Leu	Arg	Thr	Gly	Glu	Glu	Lys	Lys	Asp	Val	Ser
			405					410					415		
Ile	Ser	Arg	Leu	Arg	Glu	Ala	Phe	Ser	Leu	Arg	His	Thr	Thr	Glu	Asn

420

425

430

Lys Pro His Ser Pro Lys Thr Pro Glu Pro Arg Arg Ser Pro Leu Gly
435 440 445

Gln Lys Arg Gly Met Leu Ser Ser Ser Thr Ser Gly Ala Ile Ser Asp
450 455 460

Lys Gly Val Leu Arg Pro Gln Lys Glu Ala Val Ser Ser Ser His Gly
465 470 475 480

Pro Ser Asp Pro Thr Asp Arg Ala Glu Val Glu Lys Asp Ser Gly His
485 490 495

Gly Ser Thr Ser Val Asp Ser Glu Gly Phe Ser Ile Pro Asp Thr Gly
500 505 510

Ser His Cys Ser Ser Glu Tyr Ala Ala Ser Ser Pro Gly Asp Arg Gly
515 520 525

Ser Gln Glu His Val Asp Ser Gln Glu Lys Ala Pro Glu Thr Asp Asp
530 535 540

Ser Phe Ser Asp Val Asp Cys His Ser Asn Gln Glu Asp Thr Gly Cys
545 550 555 560

Lys Phe Arg Val Leu Pro Gln Pro Thr Asn Leu Ala Thr Pro Asn Thr
565 570 575

Lys Arg Phe Lys Lys Glu Glu Ile Leu Ser Ser Ser Asp Ile Cys Gln
580 585 590

Lys Leu Val Asn Thr Gln Asp Met Ser Ala Ser Gln Val Asp Val Ala
595 600 605

Val Lys Ile Asn Lys Lys Val Val Pro Leu Asp Phe Ser Met Ser Ser
610 615 620

Leu Ala Lys Arg Ile Lys Gln Leu His His Glu Ala Gln Gln Ser Glu
625 630 635 640

Gly Glu Gln Asn Tyr Arg Lys Phe Arg Ala Lys Ile Cys Pro Gly Glu
645 650 655

Asn Gln Ala Ala Glu Asp Glu Leu Arg Lys Glu Ile Ser Lys Thr Met
660 665 670

Phe Ala Glu Met Glu Ile Ile Gly Gln Phe Asn Leu Gly Phe Ile Ile

675

680

685

Thr Lys Leu Asn Glu Asp Ile Phe Ile Val Asp Gln His Ala Thr Asp
690 695 700

Glu Lys Tyr Asn Phe Glu Met Leu Gln Gln His Thr Val Leu Gln Gly
705 710 715 720

Gln Arg Leu Ile Ala Pro Gln Thr Leu Asn Leu Thr Ala Val Asn Glu
725 730 735

Ala Val Leu Ile Glu Asn Leu Glu Ile Phe Arg Lys Asn Gly Phe Asp
740 745 750

Phe Val Ile Asp Glu Asn Ala Pro Val Thr Glu Arg Ala Lys Leu Ile
755 760 765

Ser Leu Pro Thr Ser Lys Asn Trp Thr Phe Gly Pro Gln Asp Val Asp
770 775 780

Glu Leu Ile Phe Met Leu Ser Asp Ser Pro Gly Val Met Cys Arg Pro
785 790 795 800

Ser Arg Val Lys Gln Met Phe Ala Ser Arg Ala Cys Arg Lys Ser Val
805 810 815

Met Ile Gly Thr Ala Leu Asn Thr Ser Glu Met Lys Lys Leu Ile Thr
820 825 830

His Met Gly Glu Met Asp His Pro Trp Asn Cys Pro His Gly Arg Pro
835 840 845

Thr Met Arg His Ile Ala Asn Leu Gly Val Ile Ser Gln Asn
850 855 860

<210> 8

<211> 2771

<212> DNA

<213> Homo sapiens

<400> 8

cgaggcggat cgggtgttgc atccatggag cgagctgaga gctcgagtac agaacctgct 60
aaggccatca aacctattga tcggaagtoa gtccatcaga ttgtctctgg gcaggtggta 120
ctgagtctaa gcactgccgt aaaggagtta gtagaaaaca gtctggatgc tgggtgccact 180
aatattgac taaagcttaa ggactatgga gtggatctta ttgaagtctc agacaatgga 240
tgtggggtag aagaagaaaa cttcgaaggc ttaactctga aacatcacac atctaagatt 300
caagagtttg ccgacetaac tcaggttgaa acttttggtc ttcgggggga agotctgagc 360

tcaactttgtg	cactgagcga	tgtcaccatt	tctacctgcc	acgcacgcgc	gaagggttga	420
actcagctga	tgtttgatca	caatgggaaa	attatccaga	aaacccoccta	ccccgcoccc	480
agagggacca	cagtcagcgt	gcagcagtta	ttttccacac	tacctgtgcg	ccataaggaa	540
tttcaaagga	atattaagaa	ggagtatgcc	aaaatgggtcc	aggctcttaca	tgcatactgt	600
atcatttccg	caggcatccg	tgtaagttgc	accaatcagc	ttggacaaag	aaaacgacag	660
cctgtggtat	gcacaggttg	aagccccagc	ataaaggaaa	atatcggctc	tgtgtttggg	720
cagaagcagt	tgcaaaagcct	cattcccttt	gttcagctgc	cccctagtga	ctccgtgtgt	780
gaagagtacg	gtttgagctg	tccggatgct	ctgcataatc	ttttttacat	ctcaggtttc	840
atttcacaat	gcacgcagt	agttggaagg	agttcaacag	acagacagtt	tttctttatc	900
aaccggcgcc	cttgtgaccc	agcaaaagtc	tgcagactcg	tgaatgaggt	ctaccacatg	960
tataatgcac	accagtatcc	atttgttgtt	cttaacattt	ctgttgatct	agaatgcgtt	1020
gatatcaatg	ttaactccaga	taaaaggcaa	attttgctac	aagagggaaa	gcttttgttg	1080
gcagttttaa	agacctcttt	gataggaatg	tttgatagtg	atgtcaacaa	gctaaatgtc	1140
agtcagccga	cactgctgga	tgttgaaagt	aacttaataa	aaatgcctgc	agcggtattg	1200
gaaaagccca	tggtagaaaa	gcaggatcaa	tcccttccat	taaggactgg	agaagaaaaa	1260
aaagacgtgt	ccattttccag	actgcgagag	gccttttctc	ttcgtcacaa	aacagagaac	1320
aagcctcaca	gccccaaagc	tcagaaacca	agaaggagcc	ctctagacac	gaaaagggtg	1380
atgctgtctt	ctagcacttc	aggtgccatc	tctgacaaag	gcgtccctgag	acctcagaaa	1440
gaggcagtg	gttccagtca	cggaccocagt	gacctactcg	acagagcgga	ggtggagaag	1500
gactcggggc	acggcgagc	ttccgtggat	ctcgaggggg	tcagcatccc	agacacgggc	1560
agtcactgca	gcagcgagta	tgcggccagc	tccccagggg	acagggggctc	gcaggaaacat	1620
gtggactctc	aggagaaagg	gcctgaaact	gacgactctt	tttcagatgt	ggactgccat	1680
tcaaacccgg	aagataccgg	atgtaaat	cgagttttgc	ctcagccaac	taactctgca	1740
acccccaaaa	caaagcggtt	taaaaaagaa	gaaattcttt	ccagttctga	catttgtcaa	1800
aagttagtaa	atactcagga	catgtcagcc	tctcaggttg	atgtagctgt	gaaaattaat	1860
aagaaagtgt	tgcccttgga	cttttctatg	agttcttttag	ctaaaacgaat	aaagcaggtta	1920
catcatgaag	cacagcaaag	tgaaggggaa	cagaattaca	ggaagttag	ggcaaagatt	1980
tgtcctggag	aaaatcaagc	agccgaagat	gaactaagaa	aagagataag	taaaacgatg	2040
tttgcagaaa	tggaaatcat	tggtcagttt	aacctgggat	ttataataac	caaactgaat	2100
gaggatatct	tcatagtggg	ccagcatgcc	acggacgaga	agtataactt	cgagatgctg	2160
cagcagcaca	ccgtgtctca	ggggcagagg	ctcatagcac	ctcagactct	caacttaact	2220
gctgttaatg	aagctgttct	gatagaaaat	ctggaatat	ttagaagaag	tggctttgat	2280
tttgttatcg	atgaaaatgc	tccagtcact	gaaagggcta	aactgatttc	cttgccaact	2340
agtaaaaaact	ggacctcttg	acccaggac	gtcgatgaac	tgatcttcat	gctgagcgac	2400
agccctgggg	tcatgtgcg	gccttcccca	gtcaagcaga	tgtttgcctc	cagagcctgc	2460
cggaaagtcg	tgatgatttg	gactgctctt	aacacaagcg	agatgaagaa	actgatcacc	2520
cacatggggg	agatggacca	ccctgggaac	tgtccccatg	gaaggccaac	catgagacac	2580
atcgccaacc	tgggtgtcat	ttctcagaa	tgaccgtagt	cactgtatgt	taaatattgt	2640
tttatcgcat	atttttatgt	tttgaagac	agagtcttca	ctaacctttt	ttgttttaaa	2700
atgaaacctg	ctactttaaa	aaaataccca	tcacaccat	ttaaaagtga	tcttgagaa	2760
cttttcaaac	c					2771

<210> 9

<211> 932

<212> PRT

<213> Homo sapiens

<400> 9

Met Lys Gln Leu Pro Ala Ala Thr Val Arg Leu Leu Ser Ser Ser Gln
1 5 10 15

Ile Ile Thr Ser Val Val Ser Val Val Lys Glu Leu Ile Glu Asn Ser
20 25 30

Leu Asp Ala Gly Ala Thr Ser Val Asp Val Lys Leu Glu Asn Tyr Gly
35 40 45

Phe Asp Lys Ile Glu Val Arg Asp Asn Gly Glu Gly Ile Lys Ala Val
50 55 60

Asp Ala Pro Val Met Ala Met Lys Tyr Tyr Thr Ser Lys Ile Asn Ser
65 70 75 80

His Glu Asp Leu Glu Asn Leu Thr Thr Tyr Gly Phe Arg Gly Glu Ala
85 90 95

Leu Gly Ser Ile Cys Cys Ile Ala Glu Val Leu Ile Thr Thr Arg Thr
100 105 110

Ala Ala Asp Asn Phe Ser Thr Gln Tyr Val Leu Asp Gly Ser Gly His
115 120 125

Ile Leu Ser Gln Lys Pro Ser His Leu Gly Gln Gly Thr Thr Val Thr
130 135 140

Ala Leu Arg Leu Phe Lys Asn Leu Pro Val Arg Lys Gln Phe Tyr Ser
145 150 155 160

Thr Ala Lys Lys Cys Lys Asp Glu Ile Lys Lys Ile Gln Asp Leu Leu
165 170 175

Met Ser Phe Gly Ile Leu Lys Pro Asp Leu Arg Ile Val Phe Val His
180 185 190

Asn Lys Ala Val Ile Trp Gln Lys Ser Arg Val Ser Asp His Lys Met
195 200 205

Ala Leu Met Ser Val Leu Gly Thr Ala Val Met Asn Asn Met Glu Ser
210 215 220

Phe Gln Tyr His Ser Glu Glu Ser Gln Ile Tyr Leu Ser Gly Phe Leu
225 230 235 240

Pro Lys Cys Asp Ala Asp His Ser Phe Thr Ser Leu Ser Thr Pro Glu
245 250 255

Arg Ser Phe Ile Phe Ile Asn Ser Arg Pro Val His Gln Lys Asp Ile
 260 265 270

Leu Lys Leu Ile Arg His His Tyr Asn Leu Lys Cys Leu Lys Glu Ser
 275 280 285

Thr Arg Leu Tyr Pro Val Phe Phe Leu Lys Ile Asp Val Pro Thr Ala
 290 295 300

Asp Val Asp Val Asn Leu Thr Pro Asp Lys Ser Gln Val Leu Leu Gln
 305 310 315 320

Asn Lys Glu Ser Val Leu Ile Ala Leu Glu Asn Leu Met Thr Thr Cys
 325 330 335

Tyr Gly Pro Leu Pro Ser Thr Asn Ser Tyr Glu Asn Asn Lys Thr Asp
 340 345 350

Val Ser Ala Ala Asp Ile Val Leu Ser Lys Thr Ala Glu Thr Asp Val
 355 360 365

Leu Phe Asn Lys Val Glu Ser Ser Gly Lys Asn Tyr Ser Asn Val Asp
 370 375 380

Thr Ser Val Ile Pro Phe Gln Asn Asp Met His Asn Asp Glu Ser Gly
 385 390 395 400

Lys Asn Thr Asp Asp Cys Leu Asn His Gln Ile Ser Ile Gly Asp Phe
 405 410 415

Gly Tyr Gly His Cys Ser Ser Glu Ile Ser Asn Ile Asp Lys Asn Thr
 420 425 430

Lys Asn Ala Phe Gln Asp Ile Ser Met Ser Asn Val Ser Trp Glu Asn
 435 440 445

Ser Gln Thr Glu Tyr Ser Lys Thr Cys Phe Ile Ser Ser Val Lys His
 450 455 460

Thr Gln Ser Glu Asn Gly Asn Lys Asp His Ile Asp Glu Ser Gly Glu
 465 470 475 480

Asn Glu Glu Glu Ala Gly Leu Glu Asn Ser Ser Glu Ile Ser Ala Asp
 485 490 495

Glu Trp Ser Arg Gly Asn Ile Leu Lys Asn Ser Val Gly Glu Asn Ile
 500 505 510

Glu Pro Val Lys Ile Leu Val Pro Glu Lys Ser Leu Pro Cys Lys Val
515 520 525

Ser Asn Asn Asn Tyr Pro Ile Pro Glu Gln Met Asn Leu Asn Glu Asp
530 535 540

Ser Cys Asn Lys Lys Ser Asn Val Ile Asp Asn Lys Ser Gly Lys Val
545 550 555 560

Thr Ala Tyr Asp Leu Leu Ser Asn Arg Val Ile Lys Lys Pro Met Ser
565 570 575

Ala Ser Ala Leu Phe Val Gln Asp His Arg Pro Gln Phe Leu Ile Glu
580 585 590

Asn Pro Lys Thr Ser Leu Glu Asp Ala Thr Leu Gln Ile Glu Glu Leu
595 600 605

Trp Lys Thr Leu Ser Glu Glu Glu Lys Leu Lys Tyr Glu Glu Lys Ala
610 615 620

Thr Lys Asp Leu Glu Arg Tyr Asn Ser Gln Met Lys Arg Ala Ile Glu
625 630 635 640

Gln Glu Ser Gln Met Ser Leu Lys Asp Gly Arg Lys Lys Ile Lys Pro
645 650 655

Thr Ser Ala Trp Asn Leu Ala Gln Lys His Lys Leu Lys Thr Ser Leu
660 665 670

Ser Asn Gln Pro Lys Leu Asp Glu Leu Leu Gln Ser Gln Ile Glu Lys
675 680 685

Arg Arg Ser Gln Asn Ile Lys Met Val Gln Ile Pro Phe Ser Met Lys
690 695 700

Asn Leu Lys Ile Asn Phe Lys Lys Gln Asn Lys Val Asp Leu Glu Glu
705 710 715 720

Lys Asp Glu Pro Cys Leu Ile His Asn Leu Arg Phe Pro Asp Ala Trp
725 730 735

Leu Met Thr Ser Lys Thr Glu Val Met Leu Leu Asn Pro Tyr Arg Val
740 745 750

Glu Glu Ala Leu Leu Phe Lys Arg Leu Leu Glu Asn His Lys Leu Pro
755 760 765

Ala Glu Pro Leu Glu Lys Pro Ile Met Leu Thr Glu Ser Leu Phe Asn
770 775 780

Gly Ser His Tyr Leu Asp Val Leu Tyr Lys Met Thr Ala Asp Asp Gln
785 790 795 800

Arg Tyr Ser Gly Ser Thr Tyr Leu Ser Asp Pro Arg Leu Thr Ala Asn
805 810 815

Gly Phe Lys Ile Lys Leu Ile Pro Gly Val Ser Ile Thr Glu Asn Tyr
820 825 830

Leu Glu Ile Glu Gly Met Ala Asn Cys Leu Pro Phe Tyr Gly Val Ala
835 840 845

Asp Leu Lys Glu Ile Leu Asn Ala Ile Leu Asn Arg Asn Ala Lys Glu
850 855 860

Val Tyr Glu Cys Arg Pro Arg Lys Val Ile Ser Tyr Leu Glu Gly Glu
865 870 875 880

Ala Val Arg Leu Ser Arg Gln Leu Pro Met Tyr Leu Ser Lys Glu Asp
885 890 895

Ile Gln Asp Ile Ile Tyr Arg Met Lys His Gln Phe Gly Asn Glu Ile
900 905 910

Lys Glu Cys Val His Gly Arg Pro Phe Phe His His Leu Thr Tyr Leu
915 920 925

Pro Glu Thr Thr
930

<210> 10

<211> 3063

<212> DNA

<213> Homo sapiens

<400> 10

ggcagcagtg gctgcttgcg gctagtggat ggtaattgcc tgcctcgcgc tagcagcaag 60
ctgctctgtt aaaagcgaaa atgaaacaat tgcctgcggc aacagttcga ctcctttcaa 120
gttctcagat catcacttcg gtggtcagtg ttgtaaaaga gcttattgaa aactccttgg 180
atgctgggtg cacaagcgta gatgttaaac tggagaacta tggatttgat aaaattgagg 240
tgcagataa cggggagggt atcaaggctg ttgatgcacc tgtaatggca atgaagtact 300
acacctcaaa aataaatagt catgaagatc ttgaaaattt gacaacttac ggttttcgtg 360
gagaagcctt ggggtcaatt tgttgtatag ctgagggtttt aattacaaca agaacggctg 420

ctgataattt tagcaccag tatgttttag atggcagtg ccacatactt tctcagaaac 480
 cttcacactct tggtaacagt acaactgtaa ctgctttaag attattttaag aatctacctg 540
 taagaaagca gttttactca actgcaaaaa aatgtaaaaga tgaataaaaa aagatccaag 600
 atctctctcat gagcttttgg atccttaaac ctgacttaag gattgtcttt gtacataaca 660
 aggcagttat ttggcagaaa agcagagtat cagatcacaa gatggctctc atgtcagttc 720
 tggggactgc tgttatgaac aatatggaat cctttcagta ccactctgaa gaatctcaga 780
 tttatctcag tggatttctt ccaaatgtgt atgcagacca ctcttttact agtctttcaa 840
 caccagaaag aagttttcat ttcatataca gtgcaccagt acatcaaaaa gatattctta 900
 agttaatccg acatcattac aatctgaaat gcctaaagga atctactcgt ttgtatctgt 960
 tttttcttct gaaaatcgat gttcctacag ctgatgttga tgtataattt acaccagata 1020
 aaagccaaagt attattacaa aataaggaat ctgttttaag tgctcttgaa aatctgatga 1080
 cgacttgtaa tggaccatta cctagtacaa attcttatga aataataaaa acagatgttt 1140
 ccgcagctga catcgttctt agtaaaacag cagaaacaga tgtgcttttt aataaagtgt 1200
 aatcatctgg aaagaattat tcaaatgttg atacttcagt cattccattc caaatgata 1260
 tgcataatga tgaatctgga aaaaacactg atgattgttt aatcaccag ataagtattg 1320
 tggacttttg ttatgttcat tgtagtgtg aaatttctaa cattgataaa aacactaaga 1380
 atgcatttca ggacatttca atgagtaagt tatcatggga gaactctcag acggaatata 1440
 gtaaaacttg ttttataagt tccgttaagc acaccagctc agaaaaatgc aataaagacc 1500
 atatatagta gagtggggaa aatgaggaag aagcaggtct tgaataattt tcggaattt 1560
 ctgcagatga ttggagcagg ggaataatac ttataaaatc agtgggagag aatatgtaac 1620
 ctgtgaaat tttagtgcct gaaaaaagt taccatgtaa agtaagtaat aataattatc 1680
 caatccctga acaaatgaat cttaatgaag attcatgtaa caaaaaatc aatgtaatat 1740
 ataataaat tggaaaagtt acagcttatg atttacttag caatcgagta atcagaanaa 1800
 ccatgtcagc aagtgtcttt tttgtcag atcatcgtcc tcatgtttct atagaaaaac 1860
 ctaagactag tttagagat gcaacactac aaattgaaga actgtggaag acatttgagt 1920
 aagaggaaaa actgaaatg gaagagaagg ctactaaaga cttggaacga tacaatagtc 1980
 aaatgaagag agccattgaa caggagtcac aatgtcact aaaagatggc agaaaaaaga 2040
 taaaacccac cagcgcatg aaatttggccc agaagcacaa gttaaaaacc tcattatcta 2100
 atcaacccaa acttgatgaa ctcttcagt ccaaaattga aaaaagaagg atccataata 2160
 ttaaaatggt acagatcccc ttttctatga aaaaacttaa aataaatttt aagaaacaaa 2220
 acaaaagtga cttagaagag aaggatgaac ctgtcttgat ccacaattct aggtttctgt 2280
 atgtgtgctt aatgacatcc aaaaacagag taatgttatt aaatccatat agagttagaa 2340
 aagccctgct atttaaaaga ctctctgaga atcataaact tctctgcagag ccactggaaa 2400
 agccaattat gttaacagag agtcttttta atggatctca ttatttagag gttttatata 2460
 aaatgacagc agatgaccaa agatcacagt gatcaactta cctgtctgat ctctgtctta 2520
 cagcgaatgg tttcaagata aaattgatac caggagtttc aattactgaa aattacttgg 2580
 aaatagaagg aatggctaatt tgtctcccat tctatggagt agcagattta aaagaaattc 2640
 ttaatgctat attaaacaga aatgtaaaag aagtttatga atgtagacct cgcagaatga 2700
 taagtatttt agagggaaga gcagtgcgtc tatccagaca attaccatg tacttatcaa 2760
 aagaggacat ccaagacatt atctacagaa tgaagcacca gtttggaat gaaattaaag 2820
 agtgtgttca tggctgcoca tttttctatc atttaaccta actacatgat 2880
 taaatatgtt taagaagatt agttaccatt gaaattgggt ctgtcataaa acagcatgag 2940
 tctggtttta aattatcttt gtattattgt toacatggtt attttttaaa tgaggattca 3000
 ctgacttggt tttatattga aaaaagttcc acgtatttga gaaaacgtaa ataaactaat 3060
 aac 3063

<211> 934

<212> PRT

<213> Homo sapiens

<400> 11

Met Ala Val Gln Pro Lys Glu Thr Leu Gln Leu Glu Ser Ala Ala Glu
1 5 10 15

Val Gly Phe Val Arg Phe Phe Gln Gly Met Pro Glu Lys Pro Thr Thr
20 25 30

Thr Val Arg Leu Phe Asp Arg Gly Asp Phe Tyr Thr Ala His Gly Glu
35 40 45

Asp Ala Leu Leu Ala Ala Arg Glu Val Phe Lys Thr Gln Gly Val Ile
50 55 60

Lys Tyr Met Gly Pro Ala Gly Ala Lys Asn Leu Gln Ser Val Val Leu
65 70 75 80

Ser Lys Met Asn Phe Glu Ser Phe Val Lys Asp Leu Leu Leu Val Arg
85 90 95

Gln Tyr Arg Val Glu Val Tyr Lys Asn Arg Ala Gly Asn Lys Ala Ser
100 105 110

Lys Glu Asn Asp Trp Tyr Leu Ala Tyr Lys Ala Ser Pro Gly Asn Leu
115 120 125

Ser Gln Phe Glu Asp Ile Leu Phe Gly Asn Asn Asp Met Ser Ala Ser
130 135 140

Ile Gly Val Val Gly Val Lys Met Ser Ala Val Asp Gly Gln Arg Gln
145 150 155 160

Val Gly Val Gly Tyr Val Asp Ser Ile Gln Arg Lys Leu Gly Leu Cys
165 170 175

Glu Phe Pro Asp Asn Asp Gln Phe Ser Asn Leu Glu Ala Leu Leu Ile
180 185 190

Gln Ile Gly Pro Lys Glu Cys Val Leu Pro Gly Gly Glu Thr Ala Gly
195 200 205

Asp Met Gly Lys Leu Arg Gln Ile Ile Gln Arg Gly Gly Ile Leu Ile
210 215 220

Thr Glu Arg Lys Lys Ala Asp Phe Ser Thr Lys Asp Ile Tyr Gln Asp

225		230		235		240
Leu Asn Arg	Leu Leu Lys Gly Lys Lys Gly Glu Gln Met Asn Ser Ala					
	245			250		255
Val Leu Pro	Glu Met Glu Asn Gln Val Ala Val Ser Ser Leu Ser Ala					
	260			265		270
Val Ile Lys	Phe Leu Glu Leu Leu Ser Asp Asp Ser Asn Phe Gly Gln					
	275			280		285
Phe Glu Leu Thr Thr	Phe Asp Phe Ser Gln Tyr Met Lys Leu Asp Ile					
	290			295		300
Ala Ala Val Arg Ala	Leu Asn Leu Phe Gln Gly Ser Val Glu Asp Thr					
	305			310		315
Thr Gly Ser Gln Ser	Leu Ala Ala Leu Leu Asn Lys Cys Lys Thr Pro					
	325			330		335
Gln Gly Gln Arg Leu	Val Asn Gln Trp Ile Lys Gln Pro Leu Met Asp					
	340			345		350
Lys Asn Arg Ile Glu	Glu Arg Leu Asn Leu Val Glu Ala Phe Val Glu					
	355			360		365
Asp Ala Glu Leu Arg	Gln Thr Leu Gln Glu Asp Leu Leu Arg Arg Phe					
	370			375		380
Pro Asp Leu Asn Arg	Leu Ala Lys Lys Phe Gln Arg Gln Ala Ala Asn					
	385			390		395
Leu Gln Asp Cys Tyr	Arg Leu Tyr Gln Gly Ile Asn Gln Leu Pro Asn					
	405			410		415
Val Ile Gln Ala Leu	Glu Lys His Glu Gly Lys His Gln Lys Leu Leu					
	420			425		430
Leu Ala Val Phe Val	Thr Pro Leu Thr Asp Leu Arg Ser Asp Phe Ser					
	435			440		445
Lys Phe Gln Glu Met	Ile Glu Thr Thr Leu Asp Met Asp Gln Val Glu					
	450			455		460
Asn His Glu Phe Leu	Val Lys Pro Ser Phe Asp Pro Asn Leu Ser Glu					
	465			470		475
Leu Arg Glu Ile Met	Asn Asp Leu Glu Lys Lys Met Gln Ser Thr Leu					

485

490

495

Ile Ser Ala Ala Arg Asp Leu Gly Leu Asp Pro Gly Lys Gln Ile Lys
500 505 510

Leu Asp Ser Ser Ala Gln Phe Gly Tyr Tyr Phe Arg Val Thr Cys Lys
515 520 525

Glu Glu Lys Val Leu Arg Asn Asn Lys Asn Phe Ser Thr Val Asp Ile
530 535 540

Gln Lys Asn Gly Val Lys Phe Thr Asn Ser Lys Leu Thr Ser Leu Asn
545 550 555 560

Glu Glu Tyr Thr Lys Asn Lys Thr Glu Tyr Glu Glu Ala Gln Asp Ala
565 570 575

Ile Val Lys Glu Ile Val Asn Ile Ser Ser Gly Tyr Val Glu Pro Met
580 585 590

Gln Thr Leu Asn Asp Val Leu Ala Gln Leu Asp Ala Val Val Ser Phe
595 600 605

Ala His Val Ser Asn Gly Ala Pro Val Pro Tyr Val Arg Pro Ala Ile
610 615 620

Leu Glu Lys Gly Gln Gly Arg Ile Ile Leu Lys Ala Ser Arg His Ala
625 630 635 640

Cys Val Glu Val Gln Asp Glu Ile Ala Phe Ile Pro Asn Asp Val Tyr
645 650 655

Phe Glu Lys Asp Lys Gln Met Phe His Ile Ile Thr Gly Pro Asn Met
660 665 670

Gly Gly Lys Ser Thr Tyr Ile Arg Gln Thr Gly Val Ile Val Leu Met
675 680 685

Ala Gln Ile Gly Cys Phe Val Pro Cys Glu Ser Ala Glu Val Ser Ile
690 695 700

Val Asp Cys Ile Leu Ala Arg Val Gly Ala Gly Asp Ser Gln Leu Lys
705 710 715 720

Gly Val Ser Thr Phe Met Ala Glu Met Leu Glu Thr Ala Ser Ile Leu
725 730 735

Arg Ser Ala Thr Lys Asp Ser Leu Ile Ile Ile Asp Glu Leu Gly Arg

740

745

750

Gly Thr Ser Thr Tyr Asp Gly Phe Gly Leu Ala Trp Ala Ile Ser Glu
755 760 765

Tyr Ile Ala Thr Lys Ile Gly Ala Phe Cys Met Phe Ala Thr His Phe
770 775 780

His Glu Leu Thr Ala Leu Ala Asn Gln Ile Pro Thr Val Asn Asn Leu
785 790 795 800

His Val Thr Ala Leu Thr Thr Glu Glu Thr Leu Thr Met Leu Tyr Gln
805 810 815

Val Lys Lys Gly Val Cys Asp Gln Ser Phe Gly Ile His Val Ala Glu
820 825 830

Leu Ala Asn Phe Pro Lys His Val Ile Glu Cys Ala Lys Gln Lys Ala
835 840 845

Leu Glu Leu Glu Glu Phe Gln Tyr Ile Gly Glu Ser Gln Gly Tyr Asp
850 855 860

Ile Met Glu Pro Ala Ala Lys Lys Cys Tyr Leu Glu Arg Glu Gln Gly
865 870 875 880

Glu Lys Ile Ile Gln Glu Phe Leu Ser Lys Val Lys Gln Met Pro Phe
885 890 895

Thr Glu Met Ser Glu Glu Asn Ile Thr Ile Lys Leu Lys Gln Leu Lys
900 905 910

Ala Glu Val Ile Ala Lys Asn Asn Ser Phe Val Asn Glu Ile Ile Ser
915 920 925

Arg Ile Lys Val Thr Thr
930

<210> 12

<211> 3145

<212> DNA

<213> Homo sapiens

<400> 12

ggcgggaaac agcttagtgg gtgtgggggc gcgcattttc ttcaaccagg aggtgaggag 60
gtttcgacat ggcggtgcag ccgaaggaga cgctgcagtt ggagagcgcg gccgaggtcg 120
gcttcgtgcg cttctttcag ggcattgccg agaagccgac caccacagtg cgccttttcg 180

accggggcg	cttctatacg	gogcacggcg	aggacgcgct	gctggccggc	cgggagggtg	240
tcaagaccga	gggggtgato	aagtacatgg	ggccggcagg	agcaagaagt	ctgcagatgt	300
ttgtgcttag	taaaatgaat	tttgaatctt	ttgtaaaaga	tcttctctcg	gttcgtcagt	360
atagagttga	agttttataag	aatagagctg	gaataaaggc	atccaaaggag	aatgatgtgt	420
attttgcata	taaggcttct	cctggcaatc	tctctcagtt	tgaaagacatt	ctctttggta	480
acaatgatat	gtcagcttcc	attggtgttg	tgggtgttaa	aatgtccgca	gttgatggcc	540
agagacaggt	tggagttggg	tatgtggatt	ccatacagag	gaaactagga	ctgtgtgaat	600
tcctcgataa	tgtacagttc	tcgaatctgt	aggctctcct	catccagatt	ggaccaaggg	660
aatgtgtttt	accgggagga	gagactgctg	gagacatggg	gaaactgaga	cagataaatt	720
aaagaggaggt	aattctgata	acagaaagaa	aaaaagctga	cttttcacaa	aaagacattt	780
atocaggacct	caaccgggtg	ttgaaaggca	aaaagggaga	cgagatgaat	agtctgtaat	840
tgcagagaat	ggagaatcag	gttgacagtt	catcactgtc	tgcggtaaat	aagtttttag	900
aactctttat	agatgattcc	aactttggac	agtttgaact	gactactttt	gacttcagcc	960
agttatgtaa	tttagtgtaa	gcagcagtc	gagcccttaa	cctttttcag	actttcaga	1020
aagataccac	tgtgctctcag	tctctggctg	ccttctgtga	taagtgtaaa	accctcgaag	1080
gacaaagact	tgttaaccag	tggattaaag	agcctctcat	ggataaagaac	agaatagagg	1140
agagattgaa	tttagtggaa	gctttgttag	aagatgcaga	attgaggcag	actttcaga	1200
aagatttact	tctgtcagttc	ccagatctta	accgacttgc	caagaagttt	caaagacaag	1260
cagcaaacct	acaagattgt	taccgaactc	atcagggtat	aaatcaacta	cctaattgta	1320
tacagactgt	ggaaaataat	gaaggaaaac	accgaaat	attgtttgga	gtttttgga	1380
ctctctttac	tgtatctctg	tctgacttct	ccaagtttca	ggaaatgata	gaaacaactt	1440
tagatatgta	tcagggtggaa	aacctatgaat	tccttgtaaa	accttcattt	gatcctaatt	1500
tcagtgtaatt	aagagaataa	atgaatgact	tggaaaagaa	gatgcagta	aatcaataaa	1560
gtgcagccag	agatcttggc	ttggacctgt	gcaaacagat	taaaactggat	tccagtgcac	1620
agttttggata	ttactttctg	gtaacctgta	aggaagaaaa	agtcctctgt	aacaataaaa	1680
actttagtac	tgtagatgct	cagaagaatg	gtgttaaat	taccaacagc	aatttgaact	1740
ctttaaatga	agagtatacc	aaaaataaaa	cagaatatga	agaagccagc	gatgccattg	1800
ttaaagaaat	tgtcaatatt	tcttcaggct	atgtagaacc	aatgcagaca	ctcaatgatg	1860
tgttagctca	gctagatgct	gtgtgcagct	ttgtctcag	gtcaaatgga	gcacctgttc	1920
catatgtacg	accagccatt	ttggagaaag	gacaaggga	aattatatta	aaagcatcca	1980
ggcatgcttg	tgtttgaagt	caagatgaaa	ttgcattatt	tctcaatgac	gtatacattg	2040
aaaaagataa	acagatgttc	cacatcatta	ctggccccaa	tatggagggt	aaatcaaacat	2100
atatctgaca	aactgggggt	atagtaacta	tggcccaaat	tgggtgtttt	gtgccatgtg	2160
agtcagcaga	agtgctccatt	gtggactgca	tcttagcccg	agtaggggct	ggtagcagtc	2220
aatttgaagg	aggtctccag	ttcatggctg	aaatgttgg	aactgtctct	atcctcaggt	2280
ctgcaaccaa	agattcatta	ataatcatag	atgaattggg	aagaggaaat	tctacctacg	2340
atggatttgg	gttagcatgt	gctatatcag	aatacattgc	aacaagattt	gggtcttttt	2400
gcactgtttg	accccatatt	catgaactta	ctgcttggc	caatcagata	cttctgtgta	2460
ataatctaca	tgtcacagca	ctcaccaact	aagagacott	aactatgctt	tatcagggtga	2520
agaaaaggtgt	ctgtgatcaa	agttttggga	ttcatgttgc	agagcttgc	aatttcccta	2580
agcatgtaat	agagtgtgct	aaacagaaag	ccttggaact	taggaggttt	cagtatatgt	2640
gagaatcgca	aggatatgat	atcatggaac	cagcagcaaa	gaagtgtcat	ctggaaagag	2700
agcaaggtga	aaaaattatt	caggagttcc	tgtccaaagt	gaaacaaagt	ccctttactg	2760
aaatgtcgca	agaaaacatc	acaataaagt	taaaaacagc	aaaagctgaa	gtaatagcaa	2820
agaataatag	ctttgttaaat	gaaatcattt	cacgaataaa	agttactacg	tgaataatcc	2880
cagtaaatga	atgaaggtaa	tattgataag	ctattgtctg	taatagtttt	atatgttttt	2940
atattaaccc	tttttccata	gtgttaactg	tcagtgccca	tgggctatca	acttaataag	3000
atattttagta	atattttact	ttgaggacat	tttcaagat	ttttattttg	aaaaatgaga	3060

gctgtaactg aggactgttt gcaattgaca taggcaataa taagtgatgt gctgaatttt 3120
ataaataaaa tcatgtagtt tgtgg 3145

<210> 13
<211> 756
<212> PRT
<213> Homo sapiens

<400> 13
Met Ser Phe Val Ala Gly Val Ile Arg Arg Leu Asp Glu Thr Val Val
1 5 10 15

Asn Arg Ile Ala Ala Gly Glu Val Ile Gln Arg Pro Ala Asn Ala Ile
20 25 30

Lys Glu Met Ile Glu Asn Cys Leu Asp Ala Lys Ser Thr Ser Ile Gln
35 40 45

Val Ile Val Lys Glu Gly Gly Leu Lys Leu Ile Gln Ile Gln Asp Asn
50 55 60

Gly Thr Gly Ile Arg Lys Glu Asp Leu Asp Ile Val Cys Glu Arg Phe
65 70 75 80

Thr Thr Ser Lys Leu Gln Ser Phe Glu Asp Leu Ala Ser Ile Ser Thr
85 90 95

Tyr Gly Phe Arg Gly Glu Ala Leu Ala Ser Ile Ser His Val Ala His
100 105 110

Val Thr Ile Thr Thr Lys Thr Ala Asp Gly Lys Cys Ala Tyr Arg Ala
115 120 125

Ser Tyr Ser Asp Gly Lys Leu Lys Ala Pro Pro Lys Pro Cys Ala Gly
130 135 140

Asn Gln Gly Thr Gln Ile Thr Val Glu Asp Leu Phe Tyr Asn Ile Ala
145 150 155 160

Thr Arg Arg Lys Ala Leu Lys Asn Pro Ser Glu Glu Tyr Gly Lys Ile
165 170 175

Leu Glu Val Val Gly Arg Tyr Ser Val His Asn Ala Gly Ile Ser Phe
180 185 190

Ser Val Lys Lys Gln Gly Glu Thr Val Ala Asp Val Arg Thr Leu Pro
195 200 205

Asn Ala Ser Thr Val Asp Asn Ile Arg Ser Ile Phe Gly Asn Ala Val
210 215 220

Ser Arg Glu Leu Ile Glu Ile Gly Cys Glu Asp Lys Thr Leu Ala Phe
225 230 235 240

Lys Met Asn Gly Tyr Ile Ser Asn Ala Asn Tyr Ser Val Lys Lys Cys
245 250 255

Ile Phe Leu Leu Phe Ile Asn His Arg Leu Val Glu Ser Thr Ser Leu
260 265 270

Arg Lys Ala Ile Glu Thr Val Tyr Ala Ala Tyr Leu Pro Lys Asn Thr
275 280 285

His Pro Phe Leu Tyr Leu Ser Leu Glu Ile Ser Pro Gln Asn Val Asp
290 295 300

Val Asn Val His Pro Thr Lys His Glu Val His Phe Leu His Glu Glu
305 310 315 320

Ser Ile Leu Glu Arg Val Gln Gln His Ile Glu Ser Lys Leu Leu Gly
325 330 335

Ser Asn Ser Ser Arg Met Tyr Phe Thr Gln Thr Leu Leu Pro Gly Leu
340 345 350

Ala Gly Pro Ser Gly Glu Met Val Lys Ser Thr Thr Ser Leu Thr Ser
355 360 365

Ser Ser Thr Ser Gly Ser Ser Asp Lys Val Tyr Ala His Gln Met Val
370 375 380

Arg Thr Asp Ser Arg Glu Gln Lys Leu Asp Ala Phe Leu Gln Pro Leu
385 390 395 400

Ser Lys Pro Leu Ser Ser Gln Pro Gln Ala Ile Val Thr Glu Asp Lys
405 410 415

Thr Asp Ile Ser Ser Gly Arg Ala Arg Gln Gln Asp Glu Glu Met Leu
420 425 430

Glu Leu Pro Ala Pro Ala Glu Val Ala Ala Lys Asn Gln Ser Leu Glu
435 440 445

Gly Asp Thr Thr Lys Gly Thr Ser Glu Met Ser Glu Lys Arg Gly Pro
450 455 460

Thr Ser Ser Asn Pro Arg Lys Arg His Arg Glu Asp Ser Asp Val Glu
465 470 475 480

Met Val Glu Asp Asp Ser Arg Lys Glu Met Thr Ala Ala Cys Thr Pro
485 490 495

Arg Arg Arg Ile Ile Asn Leu Thr Ser Val Leu Ser Leu Gln Glu Glu
500 505 510

Ile Asn Glu Gln Gly His Glu Val Leu Arg Glu Met Leu His Asn His
515 520 525

Ser Phe Val Gly Cys Val Asn Pro Gln Trp Ala Leu Ala Gln His Gln
530 535 540

Thr Lys Leu Tyr Leu Leu Asn Thr Thr Lys Leu Ser Glu Glu Leu Phe
545 550 555 560

Tyr Gln Ile Leu Ile Tyr Asp Phe Ala Asn Phe Gly Val Leu Arg Leu
565 570 575

Ser Glu Pro Ala Pro Leu Phe Asp Leu Ala Met Leu Ala Leu Asp Ser
580 585 590

Pro Glu Ser Gly Trp Thr Glu Glu Asp Gly Pro Lys Glu Gly Leu Ala
595 600 605

Glu Tyr Ile Val Glu Phe Leu Lys Lys Lys Ala Glu Met Leu Ala Asp
610 615 620

Tyr Phe Ser Leu Glu Ile Asp Glu Glu Gly Asn Leu Ile Gly Leu Pro
625 630 635 640

Leu Leu Ile Asp Asn Tyr Val Pro Pro Leu Glu Gly Leu Pro Ile Phe
645 650 655

Ile Leu Arg Leu Ala Thr Glu Val Asn Trp Asp Glu Glu Lys Glu Cys
660 665 670

Phe Glu Ser Leu Ser Lys Glu Cys Ala Met Phe Tyr Ser Ile Arg Lys
675 680 685

Gln Tyr Ile Ser Glu Glu Ser Thr Leu Ser Gly Gln Gln Ser Glu Val
690 695 700

Pro Gly Ser Ile Pro Asn Ser Trp Lys Trp Thr Val Glu His Ile Val
705 710 715 720

Tyr Lys Ala Leu Arg Ser His Ile Leu Pro Pro Lys His Phe Thr Glu
725 730 735

Asp Gly Asn Ile Leu Gln Leu Ala Asn Leu Pro Asp Leu Tyr Lys Val
740 745 750

Phe Glu Arg Cys
755

<210> 14
<211> 2484
<212> DNA
<213> Homo sapiens

<400> 14
cttggctctt ctggcgccaa aatgtcgttc gtggcagggg ttattcggcg gctggacgag 60
acagtgggtg accgcatcgc ggcgggggaa gttatccagc ggccagctaa tgctatcaaa 120
gagatgattg agaactgttt agatgcaaaa tccacaagta ttcaagtgat tgttaagag 180
ggaggcctga agttgattca gatccaagac aatggcaccg ggcacaggaa agaagatctg 240
gatattgtat gtgaaaaggtt cactactagt aaactgcagt cctttgagga tttagccagt 300
atttctacct atggcctttcg aggtgaggct ttggccagca taagccatgt ggctcatgtt 360
actattacaa cgaaaacagc tgatggaaag tgtgcataca gagcaagtta ctcatatgga 420
aaactgaaag cccctcctaa accatgtgct ggcaatcaag ggacccagat cacggtggag 480
gacctttttt acaacatagc cagcaggaga aaagctttaa aaaatccaag tgaagaatat 540
gggaaaaatt tggaagtgtt tggcaggtat tcagtacaca atgcaggcat tagtttctca 600
gttaaaaaac aaggagagac agtagctgat gttaggacac tacccaatgc ctcaaccgtg 660
gacaatatto gctccatctt tggaaatgct gttagtgcag aactgataga aattggatgt 720
gaggataaaa ccttagcctt caaaatgaat gggtacatat ccaatgcaaa ctactcagt 780
aagaagtgcg tcttcttact cttcatcaac catcgtctgg tagaatcaac ttccttgaga 840
aaagccatag aaacagtgtg tgcagcctat ttgcccaaaa acacacaccc attcctgtac 900
ctcagtttag aaatcagtcg ccagaatgtg gatgttaatg tgcacccccc aaagcatgaa 960
gttcacttgc tgcacgagga gagcatcctg gagcgggtgc agcagcacat cgagagcaag 1020
cttcgtggct ccaattcttc caggatgtac ttcaccacaga ctttctacc aggaattgct 1080
ggccctctcg gggagatggt taaatccaca acaagtctga cctcgtcttc tacttctgga 1140
agtagtgata aggtctatgc ccaccagatg gttctgtacag attcccgga acagaagctt 1200
gatgcatttc tgcagcctct gagcaaaacc ctgtccagtc agccccagcg cattgtcaca 1260
gaggataaga cagatatttc tagtggcagg gctaggcagc aagatgagga gatgctgaa 1320
ctccagcccc ctgtcgaagt ggtcgccaaa aatcagagct tggaggggga tacaacaaag 1380
gggacttcag aaatgtcaga gaagagagga cctacttcca gcaaccccag aaagagacat 1440
cggaagagatt ctgatgtgga aatggtgaa gatgattccc gaaaggaagt gaactgaact 1500
tgtaccccc ggagaaggat cattaaccct actagtgttt tgagtctcca ggaagaatt 1560
aatgagcagg gacatgaggt tctccgggag atgttgata accactcctt cgtgggctgt 1620
gtgaatcctc agtgggcctt ggcacagcat caaaccaagt tataccttct caacaccacc 1680
aagcttagtg aagaactggt ctaccagata ctcatattat attttgccaa ttttgggtgt 1740
ctcaggttat cggagccagc accgtctctt gacottgcca tgctgtcctt agatagcca 1800
gagagtggct ggacagagga agatggtccc aaagaaggac ttgctgaata cattgttgag 1860

tttctgaaga agaaggctga gatgcttgca gactatttct ctttggaagt tgatgaggaa 1920
 gggaacctga ttggattacc ccttctgatt gacaactatg tgcccccttt ggagggactg 1980
 cctatcttca ttcttgcact agccactgag gtgaattggg acgaagaaaa ggaatgtttt 2040
 gaaagcctca gtaaagaatg cgctatgttc tattccatcc ggaagcagta catatctgag 2100
 gagtgcaccc tctcaggcca gcagagtga ggcctggct ccattccaaa ctctggaag 2160
 tggactgttg aacacattgt ctataaagcc ttgcgtcac acattctgcc tctaaacat 2220
 ttacacagaag atggaatat cctgcagctt gctaacctgc ctgatctata caaagtcttt 2280
 gagaggtgtt aaatatggtt atttatgcac tgtgggatgt gttcttcttt ctctgtatc 2340
 cgatacaaa gttgtgtatca aagtgtgata tacaagtgt accaacataa gtgttggtag 2400
 cacttaagac ttatacttgc ctctgtag tagtcttcta tacacagtgg attgattata 2460
 aataaataga tgtgtcttaa cata 2484

<210> 15

<211> 133

<212> PRT

<213> Homo sapiens

<400> 15

Met Glu Arg Ala Glu Ser Ser Ser Thr Glu Pro Ala Lys Ala Ile Lys
 1 5 10 15

Pro Ile Asp Arg Lys Ser Val His Gln Ile Cys Ser Gly Gln Val Val
 20 25 30

Leu Ser Leu Ser Thr Ala Val Lys Glu Leu Val Glu Asn Ser Leu Asp
 35 40 45

Ala Gly Ala Thr Asn Ile Asp Leu Lys Leu Lys Asp Tyr Gly Val Asp
 50 55 60

Leu Ile Glu Val Ser Asp Asn Gly Cys Gly Val Glu Glu Glu Asn Phe
 65 70 75 80

Glu Gly Leu Thr Leu Lys His His Thr Ser Lys Ile Gln Glu Phe Ala
 85 90 95

Asp Leu Thr Gln Val Glu Thr Phe Gly Phe Arg Gly Glu Ala Leu Ser
 100 105 110

Ser Leu Cys Ala Leu Ser Asp Val Thr Ile Ser Thr Cys His Ala Ser
 115 120 125

Ala Lys Val Gly Thr
 130

<210> 16

<211> 426
<212> DNA
<213> Homo sapiens

<400> 16
cgaggcggat cgggtgttgc atccatggag cgagctgaga gctcgagtag agaacctgct 60
aaggccatca aacctattga tcggaagtca gtccatcaga tttgctctgg gcagggtgta 120
ctgagtctaa gcactgcggt aaaggagtta gtagaaaaca gtctggatgc tgggtgccact 180
aatattgatc taaagcttaa ggactatgga gtggatctta ttgaagtttc agacaatgga 240
tgtggggtag aagaagaaaa cttcgaaggc ttaactctga aacatcacac atctaagatt 300
caagagtttg ccgacctaac tcaggttgaa acttttggct ttcgggggga agctctgagc 360
tcactttgtg cactgagcga tgtcaccatt tctacctgcc acgcatcggc gaaggttgga 420
acttga 426

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2